arXiv:1005.4712v2 [math.NT] 23 Jun 2010

THE LERCH ZETA FUNCTION I. ZETA INTEGRALS

JEFFREY C. LAGARIAS AND WEN-CHING WINNIE LI

Abstract. This is the first of four papers that study algebraic and analytic structures associated to the Lerch zeta function. This paper studies "zeta integrals" associated to the Lerch zeta function using test functions, and obtains functional equations for them. Special cases include a pair of symmetrized four-term functional equations for combinations of Lerch zeta functions, found by A. Weil, for real parameters (a, c) with $0 < a, c < 1$. It extends these functions to real a, and c, and studies limiting cases of these functions where at least one of a and c take the values 0 or 1. A main feature is that as a function of three variables (s, a, c) with a, c being real variables, the Lerch zeta function has discontinuities at integer values of a and c. For fixed s, the function $\zeta(s, a, c)$ is discontinuous on part of the boundary of the closed unit square in the (a, c)-variables, and the location and nature of these discontinuities depend on the real part $\Re(s)$ of s. Analysis of this behavior is used to determine membership of these functions in $L^p([0,1]^2, da\, dc)$ for $1 \le p < \infty$, as a function of $\Re(s)$. The paper also defines generalized Lerch zeta functions associated to the oscillator representation, and gives analogous four-term functional equations for them.

Contents

1. Introduction 1
1.1. Overview of papers 2
1.2. Background and Motivation 4
2. Main results 5
3. Analytic continuation for zeta integrals 10
4. Extended zeta integrals 16
5. Extended Lerch zeta function 19
6. Boundary cases of functional equation 22
7. Renormalized Lerch functions 28
8. Weyl algebra action 31
References 36

1. Introduction

The *Lerch zeta function*

$$\zeta(s, a, c) := \sum_{n=0}^{\infty} e^{2\pi i n a}(n + c)^{-s} \tag{1.1}$$

Date: June 20, 2010.

2000 *Mathematics Subject Classification.* Primary: 11M35.

Key words and phrases. functional equation, Hurwitz zeta function, Lerch zeta function, periodic zeta function.

The research of the first author was supported by NSF grant DMS-0500555 and DMS-0801029 and that of the second author by NSF grant DMS-0457574 and DMS-0801096.

was introduced by Lipschitz [31] in 1857, for real a and c with $c > 0$, see also Lipschitz [32]. It is named after Lerch [30], who showed in 1887 that for $\Im(a) > 0$ and $0 < c < 1$ it satisfies the three-term functional equation

$$\zeta(1-s,a,c) = (2\pi)^{-s}\Gamma(s)\left\{e^{\frac{\pi is}{2}}e^{-2\pi iac}\zeta(s,-c,a) \;+\; e^{-\frac{\pi is}{2}}e^{2\pi ic(1-a)}\zeta(s,c,1-a)\right\}\ , \tag{1.2}$$

a result which is called *Lerch's transformation formula.* cf. Erdelyi [13, p. 29]. Special cases of this function are the case $a = 0$ which gives the *Hurwitz zeta function*

$$\zeta(s,c) := \sum_{n=0}^{\infty}\frac{1}{(n+c)^s} \tag{1.3}$$

studied by Hurwitz [18], and the case $c = 1$ which gives $e^{-2\pi ia}F(a,s)$, with

$$F(a,s) := \sum_{n=1}^{\infty}\frac{e^{2\pi ina}}{n^s} \tag{1.4}$$

being the *periodic zeta function* studied in Apostol [2, p. 257]. The Riemann zeta function $\zeta(s)$ occurs as the intersection of these two special cases, i.e. when $a = 0$ and $c = 1$.

The Lerch zeta function has been extensively studied, see the books of Laurenčikas and Garunkštis [29], Srivastava and Choi [41, Chap. 2] and Kanemitsu and Terada [19, Chaps. 3-5] for many analytic results. There have been many proofs given of Lerch's transformation formula and of equivalent functional equations, see for example Apostol [1], Oberhettinger [36], Mikolas [33], Berndt [3] and Weil [43].

This is the first in a series of four papers that studies algebraic and geometric structures attached to Lerch zeta functions. We begin with a brief summary of results in this series of papers, and give a more detailed description of the main results of this paper in §2.

1.1. **Overview of papers.** In Part I our starting point is a pair of symmetric functional equations for the Lerch zeta function, viewing a and c as real variables, first given by A. Weil [43] in 1976. These involve the two functions

$$L^{\pm}(s,a,c) = \zeta(s,a,c) \pm e^{-2\pi ia}\zeta(s,1-a,1-c).$$

and relate the values of these functions at parameter values (s,a,c) to values of the same function at $(1-s,1-c,a)$. We derive functional equations for general zeta integrals of Lerch type incorporating a test function, first for values (a,c) on the unit square and then extended to all real values of (a,c). We then deduce results for $L^{\pm}(s,a,c)$ and the Lerch zeta function on the extended domain by expressing them in terms of zeta integrals. We then determine that the Lerch zeta function has discontinuities at integer values of a and c, and that the nature of these discontinuities depends on the real part of s. These discontinuities appear to be an important aspect of the behavior of this function.

In Part II ([26]) we analytically continue the Lerch zeta function $\zeta(s,a,c)$ in three complex variables to a maximal domain of holomorphy. We show it analytically continues to a multivalued function of (s, a, c) which becomes single-valued on the maximal abelian covering manifold of the manifold

$$\mathcal{M} := \{(s, a, c) \in \mathbb{C} \times (\mathbb{C} \smallsetminus \mathbb{Z}) \times (\mathbb{C} \smallsetminus \mathbb{Z})\}.$$

We determine the monodomy functions describing the multivaluedness. Positive integer values of the c-variable turn out to be removable singularities, leading to a slightly larger analytic continuation. The remaining values where either a or c are integers correspond to "singular strata" omitted from this analytic continuation. It is possible to define versions of the Lerch zeta function restricted to various "singular strata," which are *not part* of this analytic continuation. In particular $a = 0$ is a singular value, and both the Hurwitz zeta function (at $a = 0$, $c \notin \mathbb{Z}$) and the Riemann zeta function (at $a = 0, c = 1$) live on "singular strata". It remains an interesting open problem to further determine the relation of these (degenerate) "singular strata" functions to the analytic continuation above, by limiting procedures; this provides an approach to get more information on the behavior of "degenerate functions" living on the singular strata. From this perspective the discontinuity properties studied in part I reflect one aspect of the behavior of such limiting procedures.

In Part III ([27]) we make the variable change $z = e^{2\pi i a}$ and analytically continue the resulting *Lerch transcendent*

$$\Phi(s, z, c) = \sum_{n=0}^{\infty} \frac{z^n}{(n + c)^s}$$

in three complex variables. This function now lives on a covering manifold of

$$\mathcal{N} := \{(s, z, c) \in \mathbb{C} \times \left(\mathbb{P}^1(\mathbb{C}) \smallsetminus \{0, 1, \infty\}\right) \times (\mathbb{C} \smallsetminus \mathbb{Z})\}.$$

It satisfies a linear PDE over the base $\mathcal{N}$ having polynomial coefficients. This function is related to the polylogarithm under the specialization $c = 1$; here the specialization $c = 1$ lies on a regular stratum. We investigate special values of (s, a, c) where the monodromy degenerates, and determine the degeneration. Such degeneration occurs for all values $s \in \mathbb{Z}$, which emphasizes the special significance of integer values of s, some of which are "critical values" in the sense of arithmetic geometry.

In Part IV ([28]) we introduce a family of two-variable "Hecke operators" acting on the (a, c)-variables, for which the Lerch zeta function is a simultaneous eigenfunction. In suitable function spaces these satisfy the Hecke relations

$$\mathsf{T}_m \mathsf{T}_n = \mathsf{T}_n \mathsf{T}_m = \mathsf{T}_{mn}.$$

associated to $GL(1)$. We determine properties of the Hecke operator action in both the"real variables" context of part I and the "complex variables" context of parts II and III. In the real-variables context we prove a uniqueness result characterizing, for each fixed $s \in \mathbb{C}$, a two-dimensional vector space of simultaneous eigenfunctions of these operators, together with some side conditions, which generalizes a theorem of Milnor [34] for the Hurwitz zeta function. In the complex-variables context there is an additional structure, consisting of an induced action of Hecke operators on the infinite-dimensional vector space spanned by the monodromy functions. This monodromy vector space also carries an action of the fundamental group of $\mathcal{M}$ and we determine information on the commutation relations of the two-variable Hecke operators with these operators.

1.2. **Background and Motivation.** Our original objective in this work was to seek a geometric or "dynamical" interpretation for the Lerch zeta function. This was motivated by observations of the first author made in [24]. He studied two-variable dynamical zeta functions $\zeta(z, s)$, which included both an arithmetical variable s and a dynamical variable z, and observed that for certain dynamical systems (those being uniformly expansive in a suitable sense) the resulting dynamical zeta function satisfied an extra functional relation $\zeta(z, s) = \zeta(zq^{-s}, 0)$. This observation provided a framework to explain the following coincidence: the zeta function for a function field over a finite field has an arithmetical definition using the s-variable and also a dynamical definition using the z-variable, and these match under the change of variable $z = q^{-s}$. An analogous result is unknown in the number field case. The Lerch zeta function is a natural multiparameter generalization of the Riemann zeta function, and it is natural to investigate whether its extra variables might have a geometric or "dynamical" interpretation. A second objective, arising in the course of the work, was the possibility of finding a representation-theoretic interpretation of the Lerch zeta function. This possibility was based on the feature that the Lerch zeta function satisfies a linear partial differential equation in the (s, a, c)-variables which might play the role of a Laplacian, and because it is a simultaneous eigenfunction of a family of "Hecke operators," mentioned in the overview above.

These four papers taken together find various extra structures attached to the Lerch zeta function, whose form suggests the existence of a more comprehensive theory (or theories) in which they fit, addressing the objectives above. In particular, there appear to be two distinct underlying contexts in which the Lerch zeta function appears, corresponding to the "real variables" structure and the "complex variables" structure mentioned in the summary above. The real variables structure, which restricts the variables a and c to be real, imposes a twisted-periodicity condition described in this paper. The first author has since found that this structure does have a representation-theoretic interpretation cf. [25]. The complex variables structure, in which a and c are complex variables, and which leads to multivalued functions in these variables, seems "geometric" in nature. We think it will be an important problem to determine interconnecting relations between these two structures.

The results in these papers are formulated in a classical language. It seems likely that some of them may be reformulated and extended in an adelic framework. Other results seem suitable for reformulation in a framework of D-modules over complex manifolds. We hope to return to these topics in subsequent work.

It would be interesting to study if the structures studied in these papers have bearing on questions around the Riemann hypothesis. It is well known that the Riemann hypothesis fails to hold for the Lerch zeta function, but the possible validity of the weaker Lindelöf hypothesis is an open question. Garunkštis and Steuding [15] (see also

Garunkštis [14]) have put forth the proposal that the Lindelöf hypothesis may hold for the Lerch zeta function, for all real parameters (a, c). Our results in part II are compatible with this proposal, in that they imply that the Lindelöf hypothesis, if it holds for the Lerch zeta function, would then also hold for its multivalued analytic continuation, i.e. it would then hold for all branches of the function lying over real values (a, c). Concerning the Riemann hypothesis itself, the Lerch zeta function gives a multi-parameter deformation of the Riemann zeta function, in which it is located in a "singular stratum" at a (non-isolated) singular point. One may ask whether new information on the Riemann hypothesis is obtainable through taking various degenerations approaching this "singular stratum," using the extra structures available.

We conclude this discussion by noting that there are further generalizations of the Lerch zeta function twisted by Dirichlet characters $\chi \pmod{N}$. These take the form

$$L^{\pm}(\chi, s, a, c) := \sum_{n\in\mathbb{Z}} \chi(n)(\operatorname{sgn}(n+c))^k e^{2\pi i n a}(n+c)^{-s},$$

with $k = 0, 1$ and $\pm := (-1)^k$. Our results can be extended to apply to these functions, since they can be expressed as linear combinations of scaled versions of $L^{\pm}(s, a, c)$. We do not treat them in these papers, in order to reduce the notational burden.

Acknowledgments. We thank K. Prasanna and P. Sarnak for helpful comments. This work began when the first author was at AT &T Labs and the second author visited there. Both authors thank AT&T Labs for support.

2. Main results

This paper proves variants of the functional equation of the Lerch zeta function, in its four-term symmetrized form, expressed using the functions $L^{\pm}(s, a, c)$ introduced below, and extends these results to zeta integrals for other test functions. These include test functions for the oscillator representation. Our focus is then on using these zeta integrals to obtain information on the dependence of the Lerch zeta function on the a and c variables. In particular we determine continuity properties in these variables as the variables approach integer values, and membership of these functions in function spaces $L^p([0, 1]^2, dadc)$ for various values of s.

The original functional equation given by Lerch [30] in 1887 was a three-term non-symmetric functional equation, now called *Lerch's transformation formula.* In 1976 A. Weil [43, p. 57] gave a pair of symmetrized four-term functional equations, which are the ones we consider here. These two functional equations encode an invariance under the action of the additive Fourier transform on the real line.

To state the symmetrized functional equations we introduce the two functions

$$L^{\pm}(s, a, c) := \zeta(s, a, c) \pm e^{-2\pi i a}\zeta(s, 1-a, 1-c), \tag{2.1}$$

and we initially suppose the domain is $0 < a, c < 1$. For $\Re(s) > 1$ these functions have the absolutely convergent Dirichlet series representations

$$L^+(s,a,c) = \sum_{\substack{n\in\mathbb{Z}\\ n\neq -c}} e^{2\pi i n a}|n+c|^{-s}, \tag{2.2}$$

$$L^-(s,a,c) = \sum_{\substack{n\in\mathbb{Z}\\ n\neq -c}} sgn(n+c) e^{2\pi i n a}|n+c|^{-s}\ , \tag{2.3}$$

see Lemma 5.3 below.

The Weil form of the functional equations involves the completion of these functions obtained by adding appropriate gamma factors.

Theorem 2.1. (Lerch Functional Equations). *Let a, c be real with $\{(a,c) \in (0,1) \times (0,1)\}$.*

(1) The completed function

$$\hat{L}^+(s,a,c) := \pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})L^+(s,a,c) \tag{2.4}$$

analytically continues to an entire function of s, and satisfies the functional equation

$$\hat{L}^+(s,a,c) = e^{-2\pi i a c}\hat{L}^+(1-s, 1-c, a). \tag{2.5}$$

(2) The completed function

$$\hat{L}^-(s,a,c) := \pi^{-\frac{s+1}{2}}\Gamma(\frac{s+1}{2})L^-(s,a,c) \tag{2.6}$$

analytically continues to an entire function of s, and satisfies the functional equation

$$\hat{L}^-(s,a,c) = i e^{-2\pi i a c}\hat{L}^-(1-s, 1-c, a). \tag{2.7}$$

The fact that the "completed" functions $\hat{L}^{\pm}(s,a,c)$ are entire functions of s for fixed $0 < a, c < 1$ implies that for these values the (non-completed) function $L^+(s,a,c)$ necessarily has "trivial zeros" at $s = 0, -2, -4, ...$, while the (non-completed) function $L^-(s,a,c)$ has "trivial zeros" at $s = -1, -3, -5, ...$, for all values of a and c.

In §3 we derive these functional equations using a method analogous to Tate's thesis [42], assuming a, c are real with $0 < a, c < 1$. We consider a *zeta integral of Lerch type* $F_k(f; s, a, c)$ attached to a general test function $f(x)$ in the Schwartz space $\mathcal{S}(\mathbb{R})$, with $k = 0, 1$, and analytically continue these integrals to entire functions of s, for fixed (a, c) as above. The perspective of Tate's thesis has two important features which are: (i) to realize a "zeta function" as a greatest common divisor of a set of "zeta integrals" indexed by test functions; and (ii) to carry this out both in local settings and in a global adelic setting. (Compare Ramakrishnan and Valenza [39, p. 242].) Here we carry out for Lerch zeta functions the "zeta integral" step using a set of test functions at the real place which transform nicely under the Fourier transform. Our treatment is global rather than local in that it uses Poisson summation, and it emphasizes operator aspects in the use of "averaging" operators. An adelic treatment of the Lerch zeta function involves further issues, which we do not take up here.

In §4 we extend the variables in Lerch zeta integrals $F_k(f; s, a, c)$, to all values $(a, c) \in \mathbb{R}\times\mathbb{R}$. We derive a general zeta integral functional equation (Theorem 4.3) for $F_k(f; s, a, c)$, specified by a test function $f(x)$ in the Schwartz space $\mathcal{S}(\mathbb{R})$, with $k = 0, 1$. For integer values of a or c the functions are no longer entire functions, but are meromorphic functions, having possible simple poles at $s = 0$ and $s = 1$. We observe that these functions exhibit extra "twisted periodicity" relations in the (a, c)-variables.

In §5, on specializing to the Gaussian test functions $\phi_0(x) = e^{-\pi x^2}$ and $\phi_1(x) = xe^{\pi x^2}$, we obtain extensions to all $(a, c) \in \mathbb{R} \times \mathbb{R}$ of the functions $L^{\pm}(s, a, c)$ and from these define an extension $\zeta_*(s, a, c)$ of the Lerch zeta function to $(a, c) \in \mathbb{R} \times \mathbb{R}$ which preserves the symmetrized functional equations. The extended function $\zeta_*(s, a, c)$ is an entire function of s whenever both a and c are not integers, and otherwise it is meromorphic in s, with possible simple poles located only at $s = 0$ or $s = 1$. We show that for $\Re(s) > 1$ it is explicitly given for fixed (a, c) by

$$\zeta_*(s, a, c) := \sum_{n+c>0} e^{2\pi i n a}(n + c)^{-s}, \tag{2.8}$$

The extended functional equations are as follows.

Theorem 2.2. (Extended Lerch Functional Equations) *Let $(a, c) \in \mathbb{R} \times \mathbb{R}$. The extended Lerch zeta function $\zeta_*(s, a, c)$ is a meromorphic function of $s \in \mathbb{C}$, which has a simple pole at $s = 1$ if $a \in \mathbb{Z}$, and is holomorphic in $\mathbb{C}$ otherwise.*

(i) In the a-variable it is periodic, with

$$\zeta_*(s, a + 1, c) = \zeta_*(s, a, c). \tag{2.9}$$

(ii) In the c-variable it satisfies the twisted periodicity equation

$$\zeta_*(s, a, c + 1) = e^{-2\pi i a}\zeta_*(s, a, c). \tag{2.10}$$

(iii) For $k = 0$ and 1 and $\pm = (-1)^k$ the completed extended functions

$$\hat{L}_*^{\pm}(s, a, c) := \pi^{-\frac{s+k}{2}}\Gamma\left(\frac{s+k}{2}\right)\Big(\zeta_*(s, a, c) + (-1)^k e^{-2\pi i a}\zeta_*(s, 1 - a, 1 - c)\Big) \tag{2.11}$$

are meromorphic for $s \in \mathbb{C}$, and are analytic except for possible simple poles at $s = 0$ or 1. The poles occur only for $k = 0$, and then if and only if $a \in \mathbb{Z}$ or $c \in \mathbb{Z}$. The completed extended functions satisfy the functional equations, for $k = 0, 1$,

$$\hat{L}_*^{\pm}(s, a, c) = i^k e^{-2\pi i a c}\hat{L}_*^{\pm}(1 - s, 1 - c, a). \tag{2.12}$$

The extended functions $L_*^{\pm}(s, a, c)$ inherit the same symmetries as $\zeta_*(s, a, c)$; namely, for all $(a, c) \in \mathbb{R} \times \mathbb{R}$, the twisted periodicity conditions hold:

$$\begin{aligned} L_*^{\pm}(s, a + 1, c) &= L_*^{\pm}(s, a, c), \\ L_*^{\pm}(s, a, c + 1) &= e^{-2\pi i a}L_*^{\pm}(s, a, c). \end{aligned}$$

Theorem 2.2 defines the Lerch zeta function $\zeta_*(s, a, c)$ for each (a, c) on the boundary of the unit square, for all $s \in \mathbb{C}$, except for $s = 0, 1$ where poles occur. Using (2.8) and the functional equations above we find that on the boundary it is given in terms of the Hurwitz zeta function and periodic zeta function by

$$\zeta_*(s, 0, c) = \zeta(s, c) \ for\ 0 < c \le 1, \tag{2.13}$$

$$\zeta_*(s, 1, c) = \zeta(s, c) \ for\ 0 < c \le 1, \tag{2.14}$$

$$\zeta_*(s,a,0) = F(a,s) \ for \ 0 \le a \le 1, \tag{2.15}$$

and

$$\zeta_*(s,a,1) = e^{-2\pi i a} F(a,s) \ for \ 0 \le a \le 1. \tag{2.16}$$

At the four corners of the square we have

$$\zeta_*(s,0,0) = \zeta_*(s,0,1) = \zeta_*(s,1,0) = \zeta_*(s,1,1) = \zeta(s), \tag{2.17}$$

the Riemann zeta function. Theorem 2.2 therefore includes meromorphic continuations of the Hurwitz zeta function and the periodic zeta function to all $s \in \mathbb{C}$, and also gives functional equations for these functions. At the end of §5, we deduce an extended Lerch transformation formula valid for all $(a,c) \in \mathbb{R} \times \mathbb{R}$ by taking appropriate combinations of $\hat{L}_*^+(s,a,c)$ and $\hat{L}_*^-(s,a,c)$ and using suitable Gamma function identities (Theorem 5.4).

In §6 we analyze how the extended Lerch zeta function $\zeta_*(s,a,c)$ behaves as a and c approach integer values. We observe that Theorem 2.2 comes at a price: the function $\zeta_*(s,a,c)$ is *discontinuous at integer values of the c and a variables, for certain ranges of s.* Indeed the formula (2.8) reveals that for $\Re(s) > 1$ the function $\zeta_*(s,a,c)$ is discontinuous at integer values of the c-variable. It turns out to be discontinuous at integer values of the a-variable for another range of s-values. These ranges of s differ for the a-variable and the c-variable. Let $\Box = \{(a,c) : \ 0 \le a \le 1, \ 0 \le c \le 1\}$. We establish a limiting formula for $\zeta_*(s,a,c)$ with $(a,c) \in \Box^\circ$ as it approaches the boundary of $\Box$, which exhibits the discontinuities (Theorem 6.1). Using it, we obtain precise conditions when $\zeta_*(s,a,c)$ continuously extends to portions of the boundary of the unit square. Note that the equality $\zeta(s,a,c) = \zeta_*(s,a,c)$ holds for $(a,c) \in \Box^\circ$, so we may state these results for $\zeta(s,a,c)$, as follows.

Theorem 2.3. (Continuous Extension to Boundary) *For fixed $s \in \mathbb{C}$, the function $\{\zeta(s,a,c) : (a,c) \in \Box^\circ\}$ continuously extends to portions of the boundary $\partial\Box$ of the unit square to define a function $\zeta(s,a,c)$ there, in the following cases:*

(i) $a = 0$ *and* $0 < c < 1$ *when* $\Re(s) > 1$;
(ii) $a = 1$ *and* $0 < c < 1$ *when* $\Re(s) > 1$;
(iii) $c = 0$ *and* $0 < a < 1$ *when* $\Re(s) < 0$;
(iv) $c = 1$ *and* $0 < a < 1$ *when* $s \in \mathbb{C}$;
(v) *the two corners* $(a,c) = (0,1)$ *and* $(1,1)$, *when* $\Re(s) > 1$.

In these cases this continuous extension satisfies

$$\zeta(s,a,c) = \zeta_*(s,a,c). \tag{2.18}$$

A continuous extension to the boundary does not exist for any other values of (s,a,c).

The ranges where continuous extensions are possible are indicated schematically by the heavy lines in Figure 1. Interesting features to note about the discontinuities are:

(1) The locations of the discontinuities *depend only on the value of $\Re(s)$.*

(2) The locations of the discontinuities are *not invariant under the transformation* $(s,a,c) \to (1-s, 1-c, a)$ that appears in the functional equation.

(3) There is no continuous extension to the corner points $(a,c) = (0,0)$ and $(1,0)$.



FIGURE 1. Continuous extension of $\zeta(s,a,c)$ (heavy lines).

The occurrence and behavior of the discontinuities is an important feature of the Lerch zeta function. In part II we show that the integer values of a and c correspond to complex singularities to these functions, on viewing a and c as complex variables. From this perspective these discontinuities form part of the structure of these singularities. It is important to understand the nature of these singularities because the Hurwitz zeta function, Periodic zeta function and Riemann zeta function all arise in the cases where one or both a and c are integers.

In §7 we use the determination of the limiting behavior of $\zeta_*(s,a,c)$ to define "renormalized" versions $L^{R,\pm}(s,a,c)$ of the Lerch function $L^{\pm}(s,a,c)$ that continuously extend to the boundary of $\Box$, for $s \in \mathbb{C} \smallsetminus \mathbb{Z}$. The precise result is given as Theorem 7.3. The "renormalized" functions remove four terms which account for the divergent behavior at the singularity.

More generally, for fixed $s \in \mathbb{C}$, let $\mathcal{E}_s$ (for "Eigenspace") denote the complex vector space spanned by the four functions

$$\mathcal{E}_s := \mathbb{C}[L^{\pm}(s,a,c), e^{-2\pi iac}L^{\pm}(1-s,1-c,a)]. \tag{2.19}$$

The functional equation gives linear dependencies among these four functions, which show that, for each $s \in \mathbb{C}$, the vector space $\mathcal{E}_s$ is two-dimensional. In fact we have

$$\begin{aligned} \mathcal{E}_s &= \mathbb{C}[L^{\pm}(s,a,c)] \text{ for } s \in \mathbb{C} \smallsetminus \mathbb{Z}_{\leq 0} & (2.20)\\ \mathcal{E}_s &= \mathbb{C}[e^{-2\pi iac}L^{\pm}(1-s,1-c,a)] \text{ for } s \in \mathbb{C} \smallsetminus \mathbb{Z}_{\geq 1}. & (2.21) \end{aligned}$$

Note that $\zeta(s,a,c) = \frac{1}{2}\left(L^+(s,a,c) + L^-(s,a,c)\right)$ belongs to $\mathcal{E}_s$. We can now "renormalize" all functions in $\mathcal{E}_s$ using the renormalized versions of the basis functions.

We use the renormalized Lerch zeta functions to determine membership of all elements of $\mathcal{E}_s$ in the function space $L^p(\Box, da\ dc)$ for $1 \leq p \leq \infty$, as follows.

Theorem 2.4. (L^p-Membership)

(1) Suppose $1 \leq p < 2$. *Then all functions in* $\mathcal{E}_s$ *belong to* $L^p(\Box, da\ dc)$ *if and only if* $s \in \mathbb{C}$ *satisfies*

$$1 - \frac{1}{p} < \Re(s) < \frac{1}{p}.$$

Otherwise, no nonzero function in $\mathcal{E}_s$ *belongs to* $L^p(\Box, da\ dc)$.

(2) Suppose $2 \le p \le \infty$. *Then no nonzero function in* $\mathcal{E}_s$ *belongs to* $L^p(\Box, da\, dc)$, *for all* $s \in \mathbb{C}$.

The most interesting cases are $p = 1$, where $L^{\pm}(s, a, c)$ both belong to the Banach space $L^1(\Box, da\ dc)$ for s inside the critical strip $0 < \Re(s) < 1$, and $p = 2$, where no nonzero functions in $\mathcal{E}_s$ belong to the Hilbert space $L^2(\Box, da\ dc)$, for all $s \in \mathbb{C}$. Theorem 2.4 will be relevant to the Hilbert spaces studied in Part IV.

In §8 we show compatibility of the Lerch zeta function with the oscillator representation. The Schwartz space $\mathcal{S}(\mathbb{R})$ is the set of smooth vectors of a representation of (a double cover of) the group $SL(2, \mathbb{R})$, the *oscillator representation*, and also of the Heisenberg group acting on $L^2(\mathbb{R}, dx)$, a viewpoint emphasized in Howe [17]. It is closed under the action of the Fourier transform, and also closed under the action of the Weyl algebra $\mathbf{A}_1 = \mathbb{C}[x, \frac{\partial}{\partial x}]$. The Mellin transform acts as an intertwining operator, converting the Weyl algebra action into difference operators in the s-variable, and this action is also compatible with the Fourier transform. Bump and Ng [5] and Bump et al [4] observed for the Riemann zeta function that there is an an infinite family of functional equations related to the oscillator Hamiltonian. We show that this phenomenon generalizes to the Lerch zeta function, with a similar proof. We obtain an infinite family of functional equations for functions $L_n(s, a, c)$ associated to the eigenfunctions of the oscillator Hamiltonian, indexed by the nonnegative integers $n \ge 0$. The functional equations in Theorem 2.1 correspond to the cases $n = 0$ and $n = 1$, respectively, whose test functions $\phi_0(x)$ and $\phi_1(x)$ are vacuum vectors for the two irreducible subrepresentations of the oscillator representation. The result of Bump and Ng [5] for the Riemann zeta function is then in principle recoverable by specialization of variables.

Notation. The hat notation $\hat{L}$ always denotes a "completed" function multiplied by an appropriate archimedean Euler factor (gamma factor). For functions f on the real line we use $\mathcal{F}f$ to denote an additive Fourier transform and $\mathcal{M}f$ to denote a Mellin transform (multiplicative Fourier transform), as defined in §2. The variable $\epsilon \in \{-1, 1\}$ and we often write $\epsilon = (-1)^k$ for $k = 0, 1$. The function $\delta_{\mathbb{Z}}(x) = 1$ if $x \in \mathbb{Z}$ and is 0 otherwise. Here $\Re(s)$ and $\Im(s)$ denote the real and imaginary parts of a complex variable s.

3. Analytic continuation for zeta integrals

The Schwartz space $\mathcal{S}(\mathbb{R})$ consists of all smooth functions $f(x)$ such that it and all its derivatives decrease more rapidly than any power of $|x|$ as $x \to \pm\infty$. We decompose it into even and odd functions

$$\mathcal{S}(\mathbb{R}) = \mathcal{S}^{+} \oplus \mathcal{S}^{-}, \tag{3.1}$$

and write

$$f(x) = f_{+}(x) + f_{-}(x), \tag{3.2}$$

with

$$f_{+}(x) := \frac{1}{2}(f(x) + f(-x)) \quad \text{and} \quad f_{-}(x) := \frac{1}{2}(f(x) - f(-x)).$$

The *Fourier transform* $\mathcal{F}f(y)$ is given by

$$\mathcal{F}f(y) := \int_{-\infty}^{\infty} e^{-2\pi ixy} f(x)dx. \tag{3.3}$$

It takes $\mathcal{S}^+$ to $\mathcal{S}^+$, and $\mathcal{S}^-$ to $\mathcal{S}^-$, so that

$$\mathcal{F}f_\epsilon = (\mathcal{F}f)_\epsilon \qquad \text{for} \qquad \epsilon = (-1)^k, \tag{3.4}$$

and also satisfies $\mathcal{F}\mathcal{F}f(x) = f(-x)$. The functions $\phi_0(x) = e^{-\pi x^2}$ and $\phi_1(x) = xe^{-\pi x^2}$ have Fourier transforms

$$\mathcal{F}\phi_k(y) = (-i)^k \phi_k(y) \ for \ k = \ 0, \ 1. \tag{3.5}$$

We use the *two-sided Mellin transforms* $\mathcal{M}_k$ with $k = 0, 1 \pmod 2$, defined by

$$\mathcal{M}_k(f)(s) := \int_{-\infty}^{\infty} f(x)(sgn\ x)^k |x|^s \frac{dx}{|x|} \ for \ k = \ 0, \ 1. \tag{3.6}$$

The *(one-sided) Mellin transform* $\mathcal{M}$ on $\mathbb{R}^+$ is given by

$$\mathcal{M}(f)(s) := \frac{1}{2}(\mathcal{M}_0(f)(s) + \mathcal{M}_1(f)(s)) = \int_0^{\infty} f(x)x^s \frac{dx}{x}. \tag{3.7}$$

We need only consider $\mathcal{M}_0$ acting on even Schwartz functions and $\mathcal{M}_1$ acting on odd Schwartz functions, namely

$$\mathcal{M}_k(f)(s) = \mathcal{M}_k(f_\epsilon)(s) \quad \text{where} \quad \epsilon = (-1)^k, \tag{3.8}$$

since we have

$$\mathcal{M}_k(f_\epsilon)(s) \equiv 0 \quad \text{when} \quad \epsilon \ = \ (-1)^{k+1} \ . \tag{3.9}$$

The functional equations are associated to the action of certain multiplicative averaging operators on test function $f \in \mathcal{S}(\mathbb{R})$.

Definition 3.1. (1) For fixed $0 < a, \ c < 1$, and $f(x) \in \mathcal{S}(\mathbb{R})$ the (*multiplciative) averaging operator* $A^{a,c}$ is given by

$$A^{a,c}[f](x) := \sum_{n \in \mathbb{Z}} f((n+c)x)e^{2\pi ina} \ . \tag{3.10}$$

The rapid decrease of $f(x)$ at $\pm\infty$ ensures that $A^{a,c}[f](x)$ is well-defined and continuous on $\mathbb{R} \smallsetminus \{0\}$, however it may have a singularity at $x = 0$. Its Mellin transform $\mathcal{M}(A^{a,c}[f])(s)$ is well-defined for $\Re(s) > 1$.

(2) The *symmetrized (multiplicative) averaging operators* $A_k^{a,c}$ on $f \in \mathcal{S}(\mathbb{R})$are given by

$$A_k^{a,c}[f](x) := A^{a,c}[f](x) + (-1)^k e^{-2\pi ia} A^{1-a,1-c}[f](x). \tag{3.11}$$

That is,

$$\begin{aligned} A_k^{a,c}[f](x) \ &= \ \sum_{n \in \mathbb{Z}} f((n+c)x)e^{2\pi ina} \\ &\quad + (-1)^k e^{-2\pi ia} \sum_{m \in \mathbb{Z}} f((m+1-c)x)e^{-2\pi ima} \ . \end{aligned} \tag{3.12}$$

Note that

$$A^{a,c}[f](x) := \frac{1}{2}(A_0^{a,c} + A_1^{a,c})[f](x) = \sum_{n\in\mathbb{Z}} f((n+c)x)e^{2\pi ina}. \tag{3.13}$$

Our main object of study will be the Mellin transforms of functions acted on by these averaging operators.

Definition 3.2. (1) For the test function $f \in \mathcal{S}(\mathbb{R})$ and $k = 0, 1$, the *zeta integral* $F_k(f; s, a, c)$ is defined for $0 < a < 1$, $0 < c < 1$ to be the one-sided Mellin transform of $A_k^{a,c}[f]$, phase-shifted by a factor $e^{\pi iac}$, i.e. namely

$$F_k(f; s, a, c) := e^{\pi iac}\mathcal{M}(A_k^{a,c}[f])(s) = e^{\pi iac}\int_0^\infty A_k^{a,c}[f](x)x^{s-1}dx. \tag{3.14}$$

(2) The *full zeta integral* $F(f; s, a, c)$ for the test function f is

$$F(f; s, a, c) := 2e^{\pi iac}\mathcal{M}(A^{a,c}[f])(s) = F_0(f; s, a, c) + F_1(f; s, a, c). \tag{3.15}$$

These Mellin transforms are related to the Lerch zeta function as follows.

Lemma 3.3. *Let $0 < a < 1$ and $0 < c < 1$. Then for $f \in \mathcal{S}(\mathbb{R})$ and $\Re(s) > 1$ and $k = 0, 1$,*

$$\mathcal{M}(A_k^{a,c}[f])(s) = \mathcal{M}_k(f)(s)L^{\pm}(s, a, c) \tag{3.16}$$

in which $\pm = (-1)^k$ and

$$L^{\pm}(s, a, c) := \zeta(s, a, c) \pm e^{-2\pi ia}\zeta(s, 1-a, 1-c)\ . \tag{3.17}$$

Thus

$$F_k(f; s, a, c) = e^{\pi iac}\mathcal{M}_k(f)(s)L^{\pm}(s, a, c). \tag{3.18}$$

Remark. For $(a, c) \in \square^\circ$ and $\Re(s) > 1$ we have

$$L^{\pm}(s, a, c) = \sum_{n\in\mathbb{Z}} e^{2\pi ina}(sgn\,(n+c))^k|n+c|^{-s}\ . \tag{3.19}$$

Lemma 3.3 may be viewed as showing that $e^{\pi iac}L^{\pm}(s, a, c)$ is the "symbol" of the averaging operator $B_k^{a,c}$ under the Mellin transform.

Proof. We first set

$$\Lambda_k(f; s, a, c) := \mathcal{M}_k(f)(s)\left(e^{\pi iac}\zeta(s, a, c)\right).$$

The formula (3.18) is equivalent to showing that

$$F_k(f; s, a, c) = \Lambda_k(f; s, a, c) + (-1)^k e^{-\pi i(a+1-c)}\Lambda_k(f; s, 1-a, 1-c)\ . \tag{3.20}$$

To verify (3.20) we note that for $\Re(s) > 1$,

$$e^{\pi iac}\zeta(s, a, c) = \sum_{n=0}^{\infty} e^{2\pi ia(n+c/2)}(n+c)^{-s}\ ,$$

hence

$$\begin{aligned}\Lambda_k(f;s,a,c) &= \left(\sum_{n=0}^{\infty} e^{2\pi i a(n+\frac{c}{2})}(n+c)^{-s}\right)\int_{-\infty}^{\infty} f(x)(sgn(x))^k |x|^{s-1}\,dx\ . \\ &= \int_0^{\infty}\left(\sum_{n=0}^{\infty} e^{2\pi i a(n+\frac{c}{2})} f((n+c)x))\right)x^{s-1}dx \\ &\quad +(-1)^k\int_0^{\infty}\left(\sum_{n=0}^{\infty} e^{2\pi i(-a)(-n-\frac{c}{2})} f(-(n+c)x)\right)x^{s-1}dx. \qquad (3.21)\end{aligned}$$

Combined with a similar formula for $\Lambda_k(f;s,1-a,1-c)$, a detailed calculation yields

$$\begin{aligned}\Lambda_k(f;s,a,c) &+ (-1)^k e^{-\pi i(a+1-c)}\Lambda_k(f;s,1-a,1-c) \\ &= e^{2\pi i a c}\left\{\int_0^{\infty}\sum_{n\in\mathbb{Z}} e^{2\pi i n a} f((n+c)x)x^{s-1}dx\right. \\ &\quad \left. +(-1)^k e^{-2\pi i a}\int_0^{\infty}\sum_{m\in\mathbb{Z}} e^{-2\pi i m a} f((m+1-c)x)x^{s-1}dx\right\} \qquad (3.22)\end{aligned}$$

as required. ■

Lemma 3.3 shows that the dependence of $\mathcal{M}(A_k^{a,c}[f])(s)$ on the test function f is confined to the Mellin transform $\mathcal{M}_k(f)(s)$, hence we have

$$F_k(f;s,a,c) = F_k(f_\epsilon;s,a,c) \quad \text{where} \quad \epsilon = (-1)^k. \qquad (3.23)$$

Substituting (3.21) into this definition yields the integral representation

$$\begin{aligned}F_k(f;s,a,c) \ = & \quad \textstyle\int_0^{\infty}\left(\sum_{n\in\mathbb{Z}} e^{2\pi i a(n+\frac{c}{2})} f((n+c)x)\right)x^{s-1}dx \qquad (3.24)\\ +(-1)^k e^{-\pi i(a+1-c)} & \quad \textstyle\int_0^{\infty}\left(\sum_{m\in\mathbb{Z}} e^{2\pi i(1-a)(m+\frac{1-c}{2})} f((m+1-c)x)x^{s-1}dx\right).\end{aligned}$$

For even or odd Schwartz functions we have

$$F_k(f_\epsilon;s,a,c) = \begin{cases} 2\int_0^{\infty}\left(\sum_{n\in\mathbb{Z}} e^{2\pi i a(n+c/2)} f((n+c)x)\right)x^{s-1}dx & \text{if} \quad \epsilon = (-1)^k, \\ 0 & \text{if} \quad \epsilon = (-1)^{k+1}. \end{cases} \qquad (3.25)$$

This follows by pairing the terms n and $m = -n-1$ in the two integrals in (3.24): they either cancel or match according to the value of k.

We use the Poisson summation formula to obtain an analytic continuation and functional equation for $F_k(f;s,a,c)$, and, in §4, to extend the definition to $(a,c) \in \mathbb{R}\times\mathbb{R}$. This involves splitting the integral representation (3.24) into two pieces, $\int_0^1$ and $\int_1^{\infty}$. We give the $\int_1^{\infty}$ pieces the name $\Phi_k(f;s,a,c)$. If we define $\Phi(f;s,a,c)$ by

$$\Phi(f;s,a,c) := \int_1^{\infty}\left(\sum_{n\in\mathbb{Z}} e^{2\pi i a(n+c/2)} f((n+c)x)\right)x^{s-1}dx. \qquad (3.26)$$

then we have, for $k=0,1$,

$$\Phi_k(f;s,a,c) := \Phi(f;s,a,c) + (-1)^k e^{-\pi i(a+1-c)}\Phi(f;s,1-a,1-c). \qquad (3.27)$$

The function $\Phi(f; s, a, c)$ defined by (3.26) is an entire function of s on the domain $0 < a < 1, 0 < c < 1$, and the same property is inherited by $\Phi_k(f; s, a, c)$ for $k = 0, 1$.

We now show:

Lemma 3.4. *For fixed* $0 < a < 1$ *and* $0 < c < 1$ *and* $f(x) \in \mathcal{S}(\mathbb{R})$*, the function* $F_k(f; s, a, c)$ *is given by*

$$F_k(f; s, a, c) = \Phi_k(f; s, a, c) + (-1)^k e^{-\pi i a} \Phi_k(\mathcal{F}f; 1 - s, 1 - c, a) \tag{3.28}$$

for $k = 0$ *or* 1 *and* $\Re(s) > 1$*. The right side analytically continues* $F_k(f; s, a, c)$ *to an entire function of* $s \in \mathbb{C}$*.*

Proof. The integral representation (3.24) gives

$$F_k(f; s, a, c) = \Phi_k(f; s, a, c) + \Psi_k(f; s, a, c), \tag{3.29}$$

in which $\Phi_k(f; s, a, c)$ is given by (3.27) and

$$\begin{aligned}\Psi_k(f; s, a, c) := &\int_0^1 \left(\sum_{n \in \mathbb{Z}} e^{2\pi i a (n + \frac{c}{2})} f((n + c)x) \right) x^{s-1} dx \\ &+ (-1)^k e^{-\pi i (a+1-c)} \int_0^1 \left(\sum_{n \in \mathbb{Z}} e^{2\pi i (1-a)(n + \frac{1-c}{2})} f((n + 1 - c)x) \right) x^{s-1} dx\,. \end{aligned} \tag{3.30}$$

Write $f = f_+ + f_-$ as a sum of even and odd functions, and we obtain

$$\Psi_k(f_\epsilon; s, a, c) = \begin{cases} 2 \int_0^1 \left(\sum_{n \in \mathbb{Z}} e^{2\pi i a (n + \frac{c}{2})} f_\epsilon((n + c)x) \right) x^{s-1} dx & \text{if} \quad \epsilon = (-1)^k, \\ 0 & \text{if} \quad \epsilon = (-1)^{k+1}, \end{cases} \tag{3.31}$$

by the same proof as (3.25) above. Thus we have $\Psi_k(f; s, a, c) = \Psi_k(f_\epsilon; s, a, c)$ with $\epsilon = (-1)^k$. The Poisson summation formula applied to Schwartz functions gives, for $x \in \mathbb{R}_{>0}$ and any $a, c \in \mathbb{R}$,

$$\sum_{n \in \mathbb{Z}} e^{2\pi i a n} f((n + c)x) = \frac{1}{|x|} \sum_{m \in \mathbb{Z}} e^{2\pi i c (m - a)} \mathcal{F}f \left(\frac{m - a}{x} \right) . \tag{3.32}$$

Substituting this into the definition of $\Psi_k(s, a, c; f)$ yields, for $\epsilon = (-1)^k$, that

$$\Psi_k(f; s, a, c) = \Psi_k(f_\epsilon; s, a, c) = 2 \int_0^1 \left(e^{\pi i a c} \sum_{m \in \mathbb{Z}} e^{2\pi i c (m - a)} \mathcal{F}f_\epsilon \left(\frac{m - a}{x} \right) \right) x^{s-1} dx\,.$$

Making the change of variable $y = 1/x$ gives rise to

$$\Psi_k(f_\epsilon; s, a, c) = 2 \int_1^\infty \left(\sum_{m \in \mathbb{Z}} e^{2\pi i c (m - \frac{a}{2})} \mathcal{F}f_\epsilon((m - a)y) \right) y^{-s} dy\,. \tag{3.33}$$

Letting $m \to -m$ yields, since $\epsilon = (-1)^k$,

$$\begin{aligned}\Psi_k(f_\epsilon; s, a, c) &= 2\int_1^\infty \left(\sum_{m\in\mathbb{Z}} e^{2\pi ic(-m-\frac{a}{2})}\mathcal{F}f_\epsilon((-m-a)y)\right) y^{-s}dy \\ &= (-1)^k e^{-\pi ia}\left(2\int_1^\infty \sum_{m\in\mathbb{Z}} e^{2\pi i(1-c)(m+\frac{a}{2})}\mathcal{F}f_\epsilon((m+a)y)\right) y^{-s}dy \\ &= (-1)^k e^{-\pi ia}\Phi_k(\mathcal{F}f_\epsilon; 1-s, 1-c, a)\ . \qquad (3.34)\end{aligned}$$

Since $\Phi_k(f; s, a, c) = \Phi_k(f_\epsilon; s, a, c)$, by a similar proof to (3.25), we obtain from (3.34) that

$$\Psi_k(f; s, a, c) = (-1)^k e^{-\pi ia}\Phi_k(\mathcal{F}f; 1-s, 1-c, a) \qquad (3.35)$$

for all $f \in \mathcal{S}(\mathbb{R})$. Substituting this in (3.29) yields the desired result. ■

We now deduce Theorem 2.1 using Lemma 3.4 with specific test functions.

Proof of Theorem 2.1. The functional equations are associated to test functions which are self-reciprocal under the Fourier transform, in the sense that $\mathcal{F}f(x) = cf(x)$, for some constant c, necessarily a fourth root of unity. We consider the test functions

$$\phi_0(x) := e^{-\pi x^2} \in \mathcal{S}^+ \text{ and } \phi_1(x) := x\ e^{-\pi x^2} \in \mathcal{S}^-.$$

These satisfy the self-reciprocal Fourier-transform formula $\mathcal{F}\phi_0(x) = e^{-\pi x^2} = \phi_0(x)$ and $\mathcal{F}\phi_1(x) = -i\phi_1(x)$.

Suppose that $0 < a < 1$, $0 < c < 1$ and $\Re(s) > 1$. Since $\mathcal{M}_0(e^{-\pi x^2})(s) = \pi^{-s/2}\Gamma(s/2)$, the definition of $F_k(f; s, a, c)$ gives

$$\hat{L}^+(s, a, c) = \pi^{-s/2}\Gamma(s/2)(\zeta(s, a, c) + e^{-2\pi ia}\zeta(s, 1-a, 1-c)) = e^{-\pi iac}F_0(\phi_0; s, a, c)\ . \qquad (3.36)$$

Similarly since $\mathcal{M}_1(xe^{-\pi x^2})(s) = \pi^{-\frac{s+1}{2}}\Gamma\left(\frac{s+1}{2}\right)$, one finds that

$$\hat{L}^-(s, a, c) = \pi^{-\frac{s+1}{2}}\Gamma\left(\frac{s+1}{2}\right)(\zeta(s, a, c) - e^{-2\pi ia}\zeta(s, 1-a, 1-c)) = e^{-\pi iac}F_1(\phi_1; s, a, c)\ . \qquad (3.37)$$

Lemma 3.4 gives the analytic continuation of $F_k(f; s, a, c)$ to an entire function of s, which shows for $0 < a < 1, 0 < c < 1$ that each of $\hat{L}^\pm(s, a, c)$ extends to an entire function of s.

The functional equations follow from Lemma 3.4 using the definition of $\Phi_k(f; s, a, c)$ and the Fourier transform formulas. Namely

$$\begin{aligned}\hat{L}^+(s, a, c) &= e^{-\pi iac}F_0(\phi_0; s, a, c) \\ &= e^{-\pi iac}\left(\Phi_0(\phi_0; s, a, c) + e^{-\pi ia}\Phi_0(\phi_0; 1-s, 1-c, a)\right) \\ &= e^{-\pi iac}\Big(\Phi(\phi_0; s, a, c) + e^{-\pi i(a+1-c)}\Phi(\phi_0; s, 1-a, 1-c)\Big) \\ &\quad + e^{-\pi iac}\left(e^{-\pi ia}\Phi(\phi_0; 1-s, 1-c, a) + e^{-\pi ic}\Phi(\phi_0; 1-s, c, 1-a)\right),\end{aligned}$$

which agrees with

$$\begin{aligned}
e^{-2\pi iac} \quad & \hat{L}^{+}(1-s,1-c,a) \\
= \quad & e^{-2\pi iac}e^{-\pi i(1-c)a}F_0(\phi_0;1-s,1-c,a) \\
= \quad & e^{-\pi iac}e^{-\pi ia}\Big(\Phi_0(\phi_0;1-s,1-c,a)+e^{-\pi i(1-c)}\Phi_0(\phi_0;s,1-a,1-c)\Big) \\
= \quad & e^{-\pi iac}e^{-\pi ia}\Big(\Phi(\phi_0;1-s,1-c,a)+e^{-\pi i(-c-a)}\Phi(\phi_0;1-s,c,1-a)\Big) \\
& +e^{-\pi iac-\pi ia}e^{-\pi i(1-c)}\Big(\Phi(\phi_0;s,1-a,1-c)+e^{-\pi i(-a+c-1)}\Phi(\phi_0;s,a,c)\Big) \\
= \quad & e^{-\pi iac}\left(e^{-\pi ia}\Phi(\phi_0;1-s,1-c,a)+e^{-\pi ic}\Phi(\phi_0;1-s,c,1-a)\right) \\
& +e^{-\pi iac}\Big(e^{-\pi i(a+1-c)}\Phi(\phi_0;s,1-a,1-c)+\Phi(\phi_0;s,a,c)\Big).
\end{aligned}$$

A similar derivation holds for $\hat{L}^{-}(s,a,c)=ie^{-2\pi iac}\hat{L}(1-s,1-c,a)$, using $\phi_1(x)$. ∎

Remark. Weil's proof [43, p. 57] of the four term functional equations in Theorem 2.1 uses Kronecker's summation formula, which is a variant of the Poisson summation formula, applied directly to ϕ_0 and ϕ_1. The use, explicitly or implicitly, of the Poisson summation seems inevitable, in that it is known to be equivalent to the functional equation for $\zeta(s)$, see Patterson [38, Sec. 2.14].

4. Extended zeta integrals

We now extend the "zeta integrals" to apply to all $(a,c)\in\mathbb{R}\times\mathbb{R}$. Using the formulation of Lemma 3.4 for the "zeta integral" $F_k(f;s,a,c)$ that

$$F_k(f;s,a,c)=\Phi_k(f;s,a,c)+(-1)^k e^{-\pi ia}\Phi_k(\mathcal{F}f;1-s,1-c,a)$$

it suffices to extend the definition of $\Phi_k(f;s,a,c)$ to apply to all $(a,c)\in\mathbb{R}\times\mathbb{R}$. Now the function $\Phi(f;s,a,c)$ given by (3.26) is well-defined for all $(a,c)\in\mathbb{R}\times\mathbb{R}$, provided $\Re(s)<0$. When $c\notin\mathbb{Z}$, the integral representation (3.26) converges for all $s\in\mathbb{C}$ and defines $\Phi(f;s,a,c)$ as an entire function of s, and the same property is inherited by $\Phi_k(f;s,a,c)$, for $k=0,1$. In the remaining case $c\in\mathbb{Z}$ and $\Re(s)<0$, the term $n=-c$ in (3.26) contributes

$$e^{-\pi iac}f(0)\int_1^\infty x^{s-1}dx=-e^{-\pi iac}f(0)\frac{1}{s}, \tag{4.1}$$

while the remaining terms define an entire function of s. The functions $\Phi_k(f;s,a,c)$ in (3.27) have two such terms, the second with weight $(-1)^k$. Since the right side of (4.1) analytically continues to $s\in\mathbb{C}$, this motivates the following definition.

Definition 4.1. For $k = 0$ or 1, $(a, c) \in \mathbb{R} \times \mathbb{R}$ and $s \in \mathbb{C}$, define for each $f(x) \in \mathcal{S}(\mathbb{R})$,

$$\begin{aligned}\Phi_k(f; s, a, c) := & \int_1^\infty \left(\sum_{\substack{n \in \mathbb{Z} \\ n \neq -c}} e^{2\pi i a (n + \frac{c}{2})} f((n+c)x) \right) x^{s-1} dx \\ & + (-1)^k e^{-\pi i (a+1-c)} \int_1^\infty \left(\sum_{\substack{m \in \mathbb{Z} \\ m \neq c-1}} e^{2\pi i (1-a)(m + \frac{1-c}{2})} f((m+1-c)x) \right) x^{s-1} dx \\ & - \delta_{\mathbb{Z}}(c)\, e^{-\pi i a c} (1 + (-1)^k) f(0) \frac{1}{s}\ , \end{aligned} \qquad (4.2)$$

in which

$$\delta_{\mathbb{Z}}(c) = 1 \quad \text{if} \quad c \in \mathbb{Z} \quad \text{and equals} \quad 0 \quad \text{otherwise}\ . \qquad (4.3)$$

For fixed a and c, the function $\Phi(f; s, a, c)$ is a meromorphic function in s. It is holomorphic except for a possible simple pole at $s = 1$, which occurs if and only if $k = 0$, $c \in \mathbb{Z}$ and $f(0) \neq 0$. We make the following definition, which in view of Lemma 3.4 extends Definition 3.2 to $(a, c) \in \mathbb{R} \times \mathbb{R}$, and to all $s \in \mathbb{C}$.

Definition 4.2. Let $(a, c) \in \mathbb{R} \times \mathbb{R}$ and $f \in \mathcal{S}(\mathbb{R})$. Then for $k = 0,\ 1$ and all $s \in \mathbb{C}$, set

$$F_k(f; s, a, c) := \Phi_k(f; s, a, c) + (-1)^k e^{-\pi i a} \Phi_k(\mathcal{F}f; 1-s, 1-c, a). \qquad (4.4)$$

We note that the dependence of $F_k(f; s, a, c)$ on the test function f is still entirely in the Mellin transform factor $\mathcal{M}_k(f)(s)$, as in Lemma 3.3.

Theorem 4.3. (Lerch Zeta Integral Functional Equations) *Let* $(a, c) \in \mathbb{R} \times \mathbb{R}$ *and* $f(x) \in \mathcal{S}(\mathbb{R})$. *Then the functions*

$$F_k(f; s, a, c) := \Phi_k(f; s, a, c) + (-1)^k e^{-\pi i a} \Phi_k(\mathcal{F}f; 1-s, 1-c, a)$$

are meromorphic functions of s, *having the following properties.*

(i) *The function* $F_1(f; s, a, c)$ *is an entire function of* s *for all* $(a, c) \in \mathbb{R} \times \mathbb{R}$. *The function* $F_0(f; s, a, c)$ *is an entire function of* s *for* $(a, c) \in \mathbb{R} \times \mathbb{R}$ *with both* a, c *non-integers. Otherwise it is meromorphic, with possible simple poles at* $s = 0, 1$. *A pole may occur at* $s = 0$ *only if* a *is an integer, and at* $s = 1$ *only if* c *is an integer, with residues given by*

$$\begin{aligned} Res_{s=0} F_0(f; s, a, c) &= -2e^{-\pi i a c} f(0) \quad \text{if } c \in \mathbb{Z}, & (4.5) \\ Res_{s=1} F_0(f; s, a, c) &= 2e^{\pi i a c} \mathcal{F}f(0) \quad \text{if } a \in \mathbb{Z}. & (4.6) \end{aligned}$$

(ii) *The functions* $F_k(f; s, a, c)$ *are twisted periodic in the* a*-variable and* c*-variable, satisfying*

$$\begin{aligned} F_k(f; s, a+1, c) &= e^{\pi i c} F_k(f; s, a, c), & (4.7) \\ F_k(f; s, a, c+1) &= e^{-\pi i a} F_k(f; s, a, c). & (4.8) \end{aligned}$$

(iii) *The functions* $F_k(f; s, a, c)$ *satisfy the functional equations*

$$F_k(f; s, a, c) = (-1)^k e^{-\pi i a} F_k(\mathcal{F}f; 1-s, 1-c, a)\ . \qquad (4.9)$$

Proof. (i) For $(a, c) \in \mathbb{R} \times \mathbb{R}$ the definition (4.2) gives

$$\begin{aligned} F_k(f; s, a, c) &= \delta_{\mathbb{Z}}(a)\ e^{\pi iac}(1 + (-1)^k)\mathcal{F}f(0)\frac{1}{s-1} \\ &\quad -\delta_{\mathbb{Z}}(c)\ e^{-\pi iac}(1 + (-1)^k)f(0)\frac{1}{s} + R_k(f; s, a, c)\ , \end{aligned}$$

in which $R_k(f; s, a, c)$ is an entire function of s. This establishes the analytic properties of $F_k(f; s, a, c)$ in the s variable.

(ii). For $\epsilon = (-1)^k$ we have

$$\begin{aligned} \Phi_k(f_\epsilon; s, a+1, c) &= 2\int_1^\infty \left(\sum_{n\in\mathbb{Z}} e^{2\pi i(a+1)(n+\frac{c}{2})} f_\epsilon((n+c)x)\right) x^{s-1}dx \\ &= e^{\pi ic}\Phi_k(f_\epsilon; s, a, c)\ , \end{aligned} \tag{4.10}$$

and, similarly,

$$\begin{aligned} \Phi_k(f_\epsilon; s, a, c+1) &= 2\int_1^\infty \left(\sum_{n\in\mathbb{Z}} e^{2\pi ia(n+\frac{c+1}{2})} f_\epsilon((n+c+1)x)\right) x^{s-1}dx \\ &= e^{-\pi ia}\Phi_k(f_\epsilon; s, a, c)\ . \end{aligned} \tag{4.11}$$

These identities give, using (3.23),

$$\begin{aligned} F_k(f; s, a+1, c) &= F_k(f_\epsilon; s, a+1, c) \\ &= \Phi_k(f_\epsilon; s, a+1, c) + (-1)^k e^{-\pi i(a+1)}\Phi(\mathcal{F}f_\epsilon; 1-s, 1-c, a+1) \\ &= e^{\pi ic}\Phi(f_\epsilon; s, a, c) + (-1)^k e^{-\pi i(a+1)} e^{-\pi i(1-c)}\Phi(\mathcal{F}f_\epsilon; 1-s, 1-c, a) \\ &= e^{\pi ic}F_k(f_\epsilon; s, a, c) \\ &= e^{\pi ic}F_k(f; s, a, c)\ . \end{aligned}$$

In a similar fashion they also give

$$F_k(f; s, a, c+1) = e^{-\pi ia}F_k(f; s, a, c).$$

(iii) We first show that

$$\Phi_k(f; s, 1-a, 1-c) = (-1)^k e^{\pi i(a+1-c)}\Phi_k(f; s, a, c)\ . \tag{4.12}$$

We have, for $\epsilon = (-1)^k$, that

$$\begin{aligned} \Phi_k(f; s, 1-a, 1-c) &= \Phi(f_\epsilon; s, 1-a, 1-c) \\ &= 2\int_1^\infty \left(\sum_{n\in\mathbb{Z}} e^{2\pi i(1-a)(n+\frac{1-c}{2})} f_\epsilon((n+1-c)x)\right) x^{s-1}dx\,. \end{aligned}$$

Replacing n with $-n-1$ yields

$$\begin{aligned} \Phi_k(f_\epsilon; s, 1-a, 1-c) &= 2\int_1^\infty \left(\sum_{n\in\mathbb{Z}} e^{2\pi i(1-a)(-n-(\frac{1+c}{2}))} f_\epsilon((-n-c)x)\right) x^{s-1}dx \\ &= 2(-1)^k e^{\pi i(a+1-c)} \int_1^\infty \sum_{n\in\mathbb{Z}} e^{2\pi ia(n+\frac{c}{2})} f_\epsilon((n+c)x)x^{s-1}dx \\ &= (-1)^k e^{\pi i(a+1-c)}\Phi_k(f_\epsilon; s, a, c) = (-1)^k e^{\pi i(a+1-c)}\Phi_k(f; s, a, c)\ , \end{aligned}$$

which is (4.12).

We deduce the functional equation (4.9) as follows. For $\epsilon = (-1)^k$, we have

$$\begin{aligned} F_k(\mathcal{F}f; 1-s, 1-c, a) &= F_k(\mathcal{F}f_\epsilon; 1-s, 1-c, a) \\ &= \Phi_k(\mathcal{F}f_\epsilon; 1-s, 1-c, a) + (-1)^k e^{-\pi i(1-c)} \Phi_k(\mathcal{F}\mathcal{F}f_\epsilon; s, 1-a, 1-c) \\ &= (-1)^k e^{\pi i a}((-1)^k e^{-\pi i a} \Phi_k(\mathcal{F}f_\epsilon; 1-s, 1-c, a) \\ &\quad + \epsilon e^{-\pi i(a+1-c)} \Phi_k(f_\epsilon; s, 1-a, 1-c)) \\ &= (-1)^k e^{\pi i a}((-1)^k e^{-\pi i a} \Phi_k(\mathcal{F}f_\epsilon; 1-s, 1-c, a) + \Phi_k(f_\epsilon; s, a, c)) \\ &= (-1)^k e^{\pi i a} F_k(f_\epsilon; s, a, c) \\ &= (-1)^k e^{\pi i a} F_k(f; s, a, c) \ , \end{aligned}$$

in which we used $\mathcal{F}\mathcal{F}f_\epsilon(x) = f_\epsilon(-x) = \epsilon f_\epsilon(x)$, and (4.12) was used at the third step. ∎

5. Extended Lerch zeta function

We next enlarge the definition of the functions $L^{\pm}(s, a, c)$ to $(a, c) \in \mathbb{R} \times \mathbb{R}$, using certain extended zeta integrals from §4. Then we use these functions to define an extended Lerch zeta function $\zeta_*(s, a, c)$ to the same domain. We emphasize that this extended function is not obtainable by analytic continuation in the a- anc c-variables. Indeed our results show that these extended functions necessarily have discontinuities in the (a, c) variables at integer values of a and c, for some ranges of s; their virtue is that they preserve the twisted periodicity relations given in Theorem 2.2.

Using the test functions $\phi_0(x) = e^{-\pi x^2}$ and $\phi_1(x) = xe^{-\pi x^2}$ we obtained in §3 for $(a, c) \in \square^\circ$ the relations

$$\hat{L}^{\pm}(s, a, c) = e^{-\pi i a c} F_k(\phi_k; s, a, c),$$

where $k = 0, 1$ and $\pm = (-1)^k$, in (3.36), (3.37). The results of §4 now define the right side for all $(a, c) \in \mathbb{R} \times \mathbb{R}$, which motivates the following definition.

Definition 5.1. For all $(a, c) \in \mathbb{R} \times \mathbb{R}$ and all $s \in \mathbb{C}$ the (*extended*) *completed Lerch zeta functions* $L_*^{\pm}(s, a, c)$ are given by

$$\hat{L}_*^{+}(s, a, c) := e^{-\pi i a c} F_0(\phi_0; s, a, c), \tag{5.1}$$

$$\hat{L}_*^{-}(s, a, c) := e^{-\pi i a c} F_1(\phi_1; s, a, c) \ , \tag{5.2}$$

where the right hand sides are meromorphic functions of $s \in \mathbb{C}$, using Theorem 4.3.

Using this definition we reverse-engineer an extension of the Lerch zeta function $\zeta(s, a, c)$ on the domain $\square^\circ$ to a function $\zeta_*(s, a, c)$ on $(a, c) \in \mathbb{R} \times \mathbb{R}$. To do this, view the equation (2.1) as a pair of linear equations in $\zeta(s, a, c)$ and $\zeta(s, 1-a, 1-c)$, and eliminate $\zeta(s, 1-a, 1-c)$ to obtain

$$\zeta(s, a, c) = \frac{1}{2}\left(\frac{\pi^{s/2}}{\Gamma(\frac{s}{2})}\hat{L}^{+}(s, a, c) + \frac{\pi^{(s+1)/2}}{\Gamma(\frac{s+1}{2})}\hat{L}^{-}(s, a, c)\right), \tag{5.3}$$

which is valid for $0 < a < 1$, $0 < c < 1$, and $\Re(s) > 1$. The right side now extends by Definition 5.1 to all $(a, c) \in \mathbb{R} \times \mathbb{R}$, which motivates to the following definition.

Definition 5.2. For $(a, c) \in \mathbb{R} \times \mathbb{R}$ and $s \in \mathbb{C}$, the *extended Lerch zeta function* $\zeta_*(s, a, c)$ is defined by

$$\zeta_*(s,a,c) := \frac{1}{2}\left(\frac{\pi^{s/2}}{\Gamma(\frac{s}{2})}\hat{L}_*^+(s,a,c) + \frac{\pi^{(s+1)/2}}{\Gamma(\frac{s+1}{2})}\hat{L}_*^-(s,a,c)\right) \ . \tag{5.4}$$

The function $\zeta_*(s, a, c)$ is clearly meromorphic in s, and it is holomorphic in $\mathbb{C}$ except for a possible simple pole at $s = 1$ coming from $\hat{L}^+(s, a, c)$. To see this, observe that any pole at $s = 0$ of $\hat{L}^+(s, a, c)$ is cancelled by the zero of $\frac{1}{\Gamma(\frac{s}{2})}$ at $s = 0$, and the remaining terms on the right side of (5.4) are entire functions. Also (5.3) shows that for $0 < a < 1$ and $0 < c < 1$ we have

$$\zeta_*(s,a,c) = \zeta(s,a,c) \ . \tag{5.5}$$

The following lemma establishes the Dirichlet series formula (2.8) for $\zeta_*(s, a, c)$, as well as formulae for $\hat{L}^\pm(s, a, c)$.

Lemma 5.3. *For $\Re(s) > 1$ and all $(a, c) \in \mathbb{R} \times \mathbb{R}$, there holds*

$$\hat{L}_*^+(s,a,c) \ = \ \pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)\left(\sum_{\substack{n\in\mathbb{Z}\\ n\neq -c}} e^{2\pi i n a}|n+c|^{-s}\right) \ , \tag{5.6}$$

$$\hat{L}_*^-(s,a,c) \ = \ \pi^{-\frac{s+1}{2}}\Gamma\left(\frac{s+1}{2}\right)\left(\sum_{\substack{n\in\mathbb{Z}\\ n\neq -c}} sgn(n+c) e^{2\pi i n a}|n+c|^{-s}\right) \ . \tag{5.7}$$

In consequence, on the same domain,

$$\zeta_*(s,a,c) = \sum_{n+c>0} e^{2\pi i n a}(n+c)^{-s} \ . \tag{5.8}$$

Proof. For $s \in \mathbb{C}$, evaluating $\hat{L}_*^+(s, a, c)$ from its definition (5.1), using (4.4) and (4.2), yields

$$\begin{aligned}\frac{1}{2}\hat{L}_*^+(s,a,c) \ &= \ \int_1^\infty \left(\sum_{\substack{n\in\mathbb{Z}\\ n\neq -c}} e^{2\pi i a n} e^{-\pi(n+c)^2x^2}\right) x^{s-1}dx \\ &\quad + \int_1^\infty \left(\sum_{\substack{n\in\mathbb{Z}\\ n\neq a}} e^{2\pi i c(n-a)} e^{-\pi(n-a)^2y^2}\right) y^{-s}dy \\ &\quad + \delta_{\mathbb{Z}}(a)\frac{1}{s-1} - \delta_{\mathbb{Z}}(c)\ e^{-2\pi i a c}\frac{1}{s} \ .\end{aligned} \tag{5.9}$$

Since $\Re(s) > 1$ the $\delta_{\mathbb{Z}}(a)$-term may be absorbed as $n = a$ in the second integral. Applying Poisson summation (3.32) to the second integral and letting $x = \frac{1}{y}$ yields, for $\Re(s) > 1$,

$$\begin{aligned}\int_1^\infty \left(\sum_{n\in\mathbb{Z}} e^{2\pi i c(n-a)} e^{-\pi(n-a)^2y^2}\right) y^{-s}dy \ &= \ \int_0^1 \left(\sum_{\substack{n\in\mathbb{Z}\\ n\neq -c}} e^{2\pi i a n} e^{-\pi(n+c)^2x^2}\right) x^{s-1}dx \\ &\quad + \delta_{\mathbb{Z}}(c) e^{-2\pi i a c}\int_0^1 y^{s-1}dy \ .\end{aligned} \tag{5.10}$$

The last integral is $\frac{1}{s}$ when $\Re(s) > 0$, and substituting this in (5.9) yields

$$\frac{1}{2}\hat{L}_*^+(s,a,c) = \int_0^\infty \left(\sum_{\substack{n\in\mathbb{Z} \\ n\neq -c}} e^{2\pi i a n} e^{-\pi(n+c)^2x^2} \right) x^{s-1}dx \ ,$$

in which the pole terms at $s = 0$ when $c \in \mathbb{Z}$ cancel out. Since $\Re(s) > 1$ we can interchange summation with the integral, and using the change of variables $y = |n+c|x$ yields (5.6).

We can perform a similar sequence of steps on $\hat{L}_*^-(s,a,c)$ starting from

$$\begin{aligned} \frac{1}{2}\hat{L}_*^-(s,a,c) \quad &= \quad \int_1^\infty \left(\sum_{n\in\mathbb{Z}} e^{2\pi i a n}(n+c)e^{-\pi(n+c)^2x^2} \right) x^s dx \\ &\quad -i\int_1^\infty \left(\sum_{n\in\mathbb{Z}} e^{2\pi i c(n-a)}(n-a)e^{-\pi(n-a)^2y^2} \right) y^{1-s}dy, \qquad (5.11) \end{aligned}$$

which is valid for $s \in \mathbb{C}$, to obtain (5.7). Substituting these two formulas in the definition (5.4) of $\zeta_*(s,a,c)$ yields (5.8). ■

We now deduce Theorem 2.2 from Theorem 4.3 with special test functions.

Proof of Theorem 2.2. Using Lemma 5.3 we see that for $\Re(s) > 1$ the definition 5.2 for $\zeta_*(s,a,c)$ agrees with the formula (2.8),

$$\zeta_*(s,a,c) = \sum_{n+c>0} e^{2\pi i n a}(n+c)^{-s}.$$

For $\Re(s) > 1$ this function is manifestly discontinuous in the c-variable, it jumps when c is an integer. In particular, when $\Re(s) > 1$, we observe that

$$\zeta_*(s,a,c+1) = \sum_{n+c+1>0} e^{2\pi i n a}(n+c+1)^{-s} = e^{-2\pi i a} \sum_{m+c>0} e^{2\pi i m a}(m+c)^{-s},$$

which is a special case of property (ii).

Properties (i) - (iii) of Theorem 2.2 follow using definition 5.2 by simple calculations from properties (i) and (ii) of Theorem 4.3, applied to the test functions $\phi_0(x) = e^{-\pi x^2}$ and $\phi_1(x) = xe^{-\pi x^2}$. The changes in the form of the twisted periodicity (ii) and the functional equation (iii) compared to those in Theorem 4.3 come from the $e^{-\pi i a c}$ factor in $\hat{L}_*^\pm(s,a,c) = e^{-\pi i a c}F_k(\phi_k;s,a,c)$, for $k = 0, 1$, with $\pm = (-1)^k$. ■

We conclude this section by deducing Lerch's transformation formula for the extended Lerch zeta function $\zeta_*(1-s,a,c)$ from Theorem 2.2.

Theorem 5.4. (Extended Lerch's transformation formula) *For* $(a,c) \in \mathbb{R}\times\mathbb{R}$ *and all* $s \in \mathbb{C}$*,*

$$\zeta_*(1-s,a,c) = (2\pi)^{-s}\Gamma(s)\left\{ e^{\frac{\pi i s}{2}} e^{-2\pi i a c}\zeta_*(s,1-c,a) + e^{-\frac{\pi i s}{2}} e^{2\pi i c(1-a)}\zeta_*(s,c,1-a) \right\}. \tag{5.12}$$

Proof. We combine the functional equations of $\hat{L}^{\pm}(s,a,c)$ to eliminate one term. We obtain

$$\begin{aligned} 2\,\zeta_*(s,a,c) = e^{-2\pi iac}\{(&\frac{\pi^{-\frac{1-s}{2}}\Gamma(\frac{1-s}{2})}{\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})} + i\,\frac{\pi^{-\frac{2-s}{2}}\Gamma(\frac{2-s}{2})}{\pi^{-\frac{1+s}{2}}\Gamma(\frac{1+s}{2})})\zeta_*(1-s,1-c,a) \\ &+e^{2\pi ic}(\frac{\pi^{-\frac{1-s}{2}}\Gamma(\frac{1-s}{2})}{\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})} - i\,\frac{\pi^{-\frac{2-s}{2}}\Gamma(\frac{2-s}{2})}{\pi^{-\frac{1+s}{2}}\Gamma(\frac{1+s}{2})})\zeta_*(1-s,c,1-a)\}. \end{aligned} \tag{5.13}$$

Then, replacing s by $1-s$ throughout and judiciously applying the two gamma function identities

$$\Gamma(s)\Gamma(1-s) = \frac{\pi}{\sin \pi s} \tag{5.14}$$

and

$$\Gamma\left(\frac{s}{2}\right)\Gamma\left(\frac{1+s}{2}\right) = \sqrt{2\pi}2^{\frac{1}{2}-s}\Gamma(s) \tag{5.15}$$

eventually yields (5.12). ∎

6. Boundary cases of functional equation

In this section we study the behavior of $\zeta_*(s,a,c)$ as (a,c) moves from the interior of the unit square to the boundary, for fixed s; these formulas extract terms producing discontinuities at these boundaries, whose nature depends on the value of $\Re(s)$.

Theorem 6.1. (Lerch Boundary Limits) *For fixed* $s \in \mathbb{C} \setminus \mathbb{Z}$ *and* $(a,c) \in \Box^{\circ}$, *the following hold.*

(i) *For* $0 < a' < 1$ *and* $(a,c) \to (a', 1^-)$,

$$\zeta_*(s,a,c) \;\to\; \zeta_*(s,a',1). \tag{6.1}$$

(ii) *For* $0 < a' < 1$ *and* $(a,c) \to (a', 0^+)$,

$$\zeta_*(s,a,c) - c^{-s} \;\to\; \zeta_*(s,a',0). \tag{6.2}$$

(iii) *For* $0 < c' < 1$ *and* $(a,c) \to (1^-, c')$,

$$\begin{aligned} \zeta_*(s,a,c) \;\; - \;\; &\frac{1}{2}\frac{\pi^{\frac{s-1}{2}}\Gamma\left(\frac{1-s}{2}\right)}{\pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)}\left(1 - i\cot\frac{\pi s}{2}\right)e^{2\pi i(1-a)c}(1-a)^{s-1} \\ &\to\; \zeta_*(s,1,c'). \end{aligned} \tag{6.3}$$

(iv) *For* $0 < c' < 1$ *and* $(a,c) \to (0^+, c')$,

$$\zeta_*(s,a,c) - \frac{1}{2}\frac{\pi^{\frac{s-1}{2}}\Gamma\left(\frac{1-s}{2}\right)}{\pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)}\left(1 + i\cot\frac{\pi s}{2}\right)e^{-2\pi iac}a^{s-1} \;\to\; \zeta_*(s,0,c'). \tag{6.4}$$

Furthermore (i) and (ii) are valid for integer $s \leq 0$, *while (iii) and (iv) are valid for integer* $s \geq 1$.

To prove this result we begin with a preliminary lemma.

Lemma 6.2. *For $0 < t < \infty$ and $s \in \mathbb{C}$,*

$$\int_1^\infty e^{-\pi t^2 x^2} x^{s-1} dx = -\frac{1}{s} e^{-\pi t^2} + \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) t^{-s} + h(t, s)\,, \tag{6.5}$$

where $h(t,s)$ is continuous in t and meromorphic for $s \in \mathbb{C}$, with poles only at $s \in \{0, -2, -4, ...\}$, and for $s \notin 2\mathbb{Z}_{\le 0}$,

$$\lim_{t \to 0^+} h(t, s) = 0. \tag{6.6}$$

Under the same conditions

$$\int_1^\infty t e^{-\pi t^2 x^2} x^s dx = \pi^{-\frac{s+1}{2}} \Gamma\left(\frac{s+1}{2}\right) t^{-s} + k(t, s), \tag{6.7}$$

where $k(t,s)$ is continuous in t and meromorphic in $s \in \mathbb{C}$, with poles only at $s \in \{-1, -3, -5, ...\}$, and for $s \notin -1 + 2\mathbb{Z}_{\le 0}$,

$$\lim_{t \to 0^+} k(t, s) = 0. \tag{6.8}$$

Proof. It suffices to prove the statement in (6.6) for $s \in \mathbb{C} \smallsetminus 2\mathbb{Z}_{\le 0}$ since both sides of (6.5) are meromorphic in s. Let $m \in \mathbb{Z}_{\ge 0}$ be such that $\Re(s + 2m) > 0$. Integrating by parts repeatedly gives

$$\begin{aligned}
\int_1^\infty e^{-\pi t^2 x^2} x^{s-1} dx &= e^{-\pi t^2 x^2} \frac{x^s}{s}\Big|_1^\infty + \frac{2\pi t^2}{s} \int_1^\infty e^{-\pi t^2 x^2} x^{s+1} dx \\
&= -\frac{e^{-\pi t^2}}{s} + \frac{2\pi t^2}{s} \int_1^\infty e^{-\pi t^2 x^2} x^{s+1} dx \\
&= -\sum_{k=0}^{m} \frac{(2\pi)^k t^{2k} e^{-\pi t^2}}{s(s+2)\cdots(s+2k)} \\
&\quad + \frac{(2\pi)^{m+1} t^{2m+2}}{s(s+2)\cdots(s+2m)} \int_1^\infty e^{-\pi t^2 x^2} x^{s+2m+1} dx\,.
\end{aligned}$$

Now write the last integral as $\int_0^\infty - \int_0^1$ and note that $\int_0^1 e^{-\pi t^2 x^2} x^{s+2m+1} dx$ converges absolutely since $\Re(s + 2m + 1) > 0$. We obtain

$$\int_1^\infty e^{-\pi t^2 x^2} x^{s-1} dx = -\frac{1}{s} e^{-\pi t^2} + h(t, s) + \frac{(2\pi)^{m+1} t^{2m+2}}{s(s+2)\cdots(s+2m)} \int_0^\infty e^{-x t^2 x^2} x^{s+2m+1} dx\,, \tag{6.9}$$

where we define

$$h(t, s) := -\sum_{j=1}^{m} \frac{(2\pi)^j t^{2j} e^{-\pi t^2}}{s(s+2)\cdots(s+2j)} - \frac{(2\pi)^{m+1} t^{2m+2}}{s(s+2)\cdots(s+2m)} \int_0^1 e^{-\pi t^2 x^2} x^{s+2m+1} dx\,. \tag{6.10}$$

Using the change of variable $y = tx$, we can rewrite the last term in (6.9) as

$$\begin{aligned}
\frac{(2\pi)^{m+1} t^{2m+2}}{s(s+2)\cdots(s+2m)} \int_0^\infty e^{-\pi t^2 x^2} x^{s+2m+1} dx &= \frac{\pi^{m+1} t^{-s}}{\frac{s}{2}\left(\frac{s}{2}+1\right)\cdots\left(\frac{s}{2}+m\right)} \int_0^\infty e^{-\pi y^2} y^{s+2m+2} \frac{dy}{y} \\
&= \frac{\pi^{m+1} t^{-s} \pi^{-\frac{s+2m+2}{2}} \Gamma\left(\frac{s+2m+2}{2}\right)}{\frac{s}{2}\left(\frac{s}{2}+1\right)\cdots\left(\frac{s}{2}+m\right)} \\
&= \pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) t^{-s}\,,
\end{aligned} \tag{6.11}$$

where we repeatedly used $s\Gamma(s) = \Gamma(s+1)$. The definition (6.10) is valid for $0 \le t < \infty$ and defines $h(t,s)$ as a meromorphic function for $\Re(s+2m+1) > 0$, and (6.11) shows that the definition of $h(t,s)$ is independent of m. It follows that $h(t,s)$ is meromorphic for $s \in \mathbb{C}$ with possible poles at $s \in 2\mathbb{Z}_{\le 0}$. For fixed $s \notin 2\mathbb{Z}_{\le 0}$ it is continuous in t, and satisfies

$$h(0,s) \equiv 0 \ . \tag{6.12}$$

Now (6.5) and $\lim_{t\to 0^+} h(t,s) = 0$ follow from (6.9)–(6.12).

We obtain the second formula (6.7) from (6.5) by multiplying by t, replacing s with $s+1$ and setting

$$k(t,s) := -\frac{te^{-\pi t^2}}{s+1} + t\ h(t,s+1)$$

as the remainder term in (6.7). This function is continuous in t for $0 \le t < \infty$ and $k(0,s) \equiv 0$, except possibly for $s \in 1 + 2\mathbb{Z}_{\le -1}$. ∎

Proof of Theorem 6.1. We define

$$\begin{aligned} \hat{L}_0^T(s,a,c) &:= \int_1^\infty \left(\sum_{\substack{n\in\mathbb{Z} \\ n\neq 0,-1}} e^{2\pi i a n} e^{-\pi(n+c)^2x^2} \right) x^{s-1} dx \\ &\quad + \int_1^\infty \left(\sum_{\substack{n\in\mathbb{Z} \\ n\neq 0,1}} e^{2\pi i c(n-a)} e^{-\pi(n-a)^2y^2} \right) y^{-s} dy \end{aligned} \tag{6.13}$$

and

$$\begin{aligned} \hat{L}_1^T(s,a,c) &:= \int_1^\infty \sum_{\substack{n\in\mathbb{Z} \\ n\neq 0,-1}} e^{2\pi i a n}(n+c) e^{-\pi(n+c)^2x^2} x^{s} dx \\ &\quad -i \int_1^\infty \left(\sum_{\substack{n\in\mathbb{Z} \\ n\neq 0,1}} e^{2\pi i c(n-a)}(n-a) e^{-\pi(n-a)^2y^2} \right) y^{1-s} dy \, . \end{aligned} \tag{6.14}$$

It is easy to see that $\hat{L}_0^T(s,a,c)$ and $\hat{L}_1^T(s,a,c)$ are entire functions in s and are jointly continuous in (a,c) for $(a,c) \in (-1,2)\times(-1,2)$. For $0 < a < 1$ and $0 < c < 1$ we have, by definition,

$$\begin{aligned} \hat{L}^+(s,a,c) &= \hat{L}_0^T(s,a,c) + \int_1^\infty e^{-\pi c^2x^2} x^{s-1} dx + e^{-2\pi i a} \int_1^\infty e^{-\pi(c-1)^2x^2} x^{s-1} dx \\ &+ \int_1^\infty e^{-2\pi i a c} e^{-\pi a^2 y^2} y^{-s} dy + \int_1^\infty e^{2\pi i c(1-a)} e^{-\pi(1-a)^2y^2} y^{-s} dy, \end{aligned} \tag{6.15}$$

and

$$\begin{aligned} \hat{L}^-(s,a,c) &= \hat{L}_1^T(s,a,c) + \int_1^\infty c e^{-\pi c^2x^2} x^{s} dx + \int_1^\infty e^{-2\pi i a}(c-1) e^{-\pi(c-1)^2x^2} x^{s} dx \\ &-i \int_1^\infty e^{-2\pi i a c}(-a) e^{-\pi a^2 y^2} y^{1-s} dy - i \int_1^\infty e^{2\pi i c(1-a)}(1-a) e^{-\pi(1-a)^2y^2} y^{1-s} dy \end{aligned} \tag{6.16}$$

We evaluate $\zeta_*(s,a,c)$ using the formula

$$\zeta_*(s,a,c) = \frac{1}{2}\left(\frac{\hat{L}^+(s,a,c)}{\pi^{-\frac{s}{2}}\Gamma(s/2)} + \frac{\hat{L}^-(s,a,c)}{\pi^{-\frac{1+s}{2}}\Gamma\left(\frac{1+s}{2}\right)}\right). \tag{6.17}$$

When considering the limit of $\hat{L}^{\pm}(s,a,c)$ as (a,c) approaches a point (a',c') on the boundary of $\square$, only those integrals on the right side of (6.15) (resp. (6.16)) in which the exponential function has exponent approaching 0 can introduce discontinuities. We call these the *bad terms*. All the remaining terms converge to the corresponding terms in $\hat{L}^{\pm}(s,a',c')$ in (5.9) and are represented by the corresponding integrals. At the point (a',c') the bad term in $\zeta_*(s,a',c')$ is replaced by the quantity

$$\delta_{\mathbb{Z}}(a')\frac{1}{s-1} - \delta_{\mathbb{Z}}(c')\frac{e^{-2\pi i a'c'}}{s}. \tag{6.18}$$

The problem is that for certain s the bad terms do not always continuously approach these values as $(a,c)\to(a',c')$.

Case I (c=1). $(a',c')=(a',1)$ with $0<a'<1$.

There is exactly one bad term in $\hat{L}^+(s,a,c)$, namely $e^{-2\pi ia}\int_1^\infty e^{-\pi(1-c)^2x^2}x^{s-1}dx$. For it Lemma 6.2 gives, for $s\in\mathbb{C}\smallsetminus 2\mathbb{Z}_{\le 0}$,

$$e^{-2\pi ia}\int_1^\infty e^{-\pi(1-c)^2x^2}x^{s-1}dx = -e^{-2\pi ia}\frac{e^{-\pi(1-c)^2}}{s}+\pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)e^{-2\pi ia}(1-c)^{-s}+e^{-2\pi ia}h(1-c,s), \tag{6.19}$$

with $h(1-c,s)\to 0$ as $c\to 0$. As $(a,c)\to(a',1)$, the first term on the right side converges to $\frac{-e^{-2\pi ia'}}{s}$, and the last term converges to zero.

There is one bad term in $\hat{L}^-(s,a,c)$, and for it Lemma 6.2 gives, for $s\in\mathbb{C}\smallsetminus\{-1+2\mathbb{Z}_{<0}\}$,

$$e^{-2\pi ia}\int_1^\infty (c-1)e^{-\pi(c-1)^2x^2}x^{s}dx = -\pi^{-\frac{s+1}{2}}\Gamma\left(\frac{s+1}{2}\right)e^{-2\pi ia}(1-c)^{-s}+e^{-2\pi ia}k(1-c,s), \tag{6.20}$$

where $k(1-c,s)\to 0$ as $c\to 1$. When these formulas are substituted in (6.17), the middle term on the right side of (6.19) cancels with the gamma-factor term on the right side of (6.20). This yields (6.1), when $s\in\mathbb{C}\smallsetminus\mathbb{Z}_{\le 0}$.

Case II. (c=0). $(a',c')=(a',0)$ with $0<a'<1$.

Here $\hat{L}^+(s,a,c)$ has a single bad term, for which Lemma 6.2 gives, for $s\in\mathbb{C}\smallsetminus 2\mathbb{Z}_{\le 0}$,

$$\int_1^\infty e^{-\pi c^2x^2}x^{s-1}dx = -\frac{e^{-\pi c^2}}{s}+\pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)c^{-s}+h(c,s). \tag{6.21}$$

As $(a,c)\to(a',0)$ the first term on the right converges to the constant $-\frac{1}{s}$, and the last term converges to zero. The middle term, when substituted in (6.17), contributes $\frac{1}{2}c^{-s}$ to the right side of (6.17).

Now $\hat{L}^-(s,a,c)$ has a single bad term, and a similar analysis shows, for $s\in\mathbb{C}\smallsetminus\{-1+2\mathbb{Z}_{\le 0}\}$, that it also contributes $\frac{1}{2}c^{-s}$ to the right side of (6.17). Put together, we obtain as $(a,c) \to (a',0^+)$,

$$\begin{aligned}\zeta_*(s,a,c) - c^{-s} &= \frac{1}{2}\left(\frac{\hat{L}^+(s,a,c)}{\pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)} - c^{-s}\right) + \frac{1}{2}\left(\frac{\hat{L}^-(s,a,c)}{\pi^{-\frac{s+1}{2}}\Gamma\left(\frac{s+1}{2}\right)} - c^{-s}\right) \\ &\to \zeta_*(s,a',0),\end{aligned}$$

which is (6.2), when $s \in \mathbb{C} \smallsetminus \mathbb{Z}_{\le 0}$.

Case III (a=1). $(a',c') = (1,c')$, with $0 < c' < 1$.

There is one bad term in $\hat{L}^+(s,a,c)$, which using Lemma 6.2 is, for $1-s \in \mathbb{C} \smallsetminus 2\mathbb{Z}_{\le 0}$,

$$\begin{aligned}&\int_1^\infty e^{2\pi ic(1-a)} e^{-\pi(1-a)^2y^2} y^{-s}dy = -e^{2\pi ic(1-a)}\frac{e^{-\pi(1-a)^2}}{1-s} \\ &+\pi^{-\frac{1-s}{2}}\Gamma\left(\frac{1-s}{2}\right) e^{2\pi ic(1-a)}(1-a)^{s-1} + e^{2\pi ic(1-a)}h(1-a,1-s)\,.\end{aligned}$$

As $(a,c) \to (1,c')$ the first term on the right side converges to the constant $\frac{1}{s-1}$, and the last term vanishes. There is one bad term in $\hat{L}^-(s,a,c)$, which using Lemma 6.2 is, for $1-s \in \mathbb{C} \smallsetminus \{-1+2\mathbb{Z}_{\le 0}\}$,

$$\begin{aligned}-i\int_1^\infty e^{2\pi ic(1-a)}(1-a)e^{-\pi(1-a)^2y^2}y^{1-s}dy = &- i\pi^{-\frac{2-s}{2}}\Gamma\left(\frac{2-s}{2}\right)e^{2\pi ic(1-a)}(1-a)^{s-1} \\ &- ie^{2\pi ic(1-a)}k(1-a,1-s)\,,\end{aligned}$$

in which the second term on the right side vanishes as $(a,c) \to (1,c')$. Substituting these into (6.17), and subtracting off the remaining two terms of the right sides yields, as $(a,c) \to (1,c')$,

$$\zeta_*(s,a,c) - \gamma(s)e^{2\pi ic(1-a)}(1-a)^{s-1} \to \zeta_*(s,1,c'), \tag{6.22}$$

in which

$$\gamma(s) := \frac{1}{2}\left(\frac{\pi^{-\frac{1-s}{2}}\Gamma\left(\frac{1-s}{2}\right)}{\pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)} - i\frac{\pi^{-\frac{2-s}{2}}\Gamma\left(\frac{2-s}{2}\right)}{\pi^{-\frac{s+1}{2}}\Gamma\left(\frac{s+1}{2}\right)}\right). \tag{6.23}$$

Using the identity $\frac{\Gamma(\frac{2-s}{2})}{\Gamma(\frac{s+1}{2})} = \frac{\Gamma(\frac{1-s}{2})}{\Gamma(\frac{s}{2})}\cot\frac{\pi s}{2}$, the formulas (6.22) and (6.23) yield (6.3), for $s \in \mathbb{C} \smallsetminus \mathbb{Z}_{\ge 1}$.

Case IV (a=0). $(a',c') = (0,c')$, with $0 < c' < 1$.

This case is parallel to case III. ∎

It is clear from the proof above that the case when (a,c) tends to a corner of the unit square can be handled in a similar fashion, except that each of $\hat{L}^+(s,a,c)$ and $\hat{L}^-(s,a,c)$ now have two bad terms which must be taken into account. The result for this case is as follows.

Theorem 6.3. (Lerch Corner Limits) *For fixed $s \in \mathbb{C} \smallsetminus \mathbb{Z}$ and $(a,c) \in \Box^\circ$ the following hold.*

(i) *As $(a,c) \to (1^-,1^-)$,*

$$\zeta_*(s,a,c) - \frac{1}{2}\frac{\pi^{-\frac{1-s}{2}}\Gamma(\frac{1-s}{2})}{\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})}\left(1 - i\cot\frac{\pi s}{2}\right)e^{2\pi i(1-a)c}(1-a)^{s-1} \to \zeta_*(s,1,1)\,.$$

(ii) *As* $(a,c) \to (0^+, 1^-)$,

$$\zeta_*(s,a,c) \quad - \quad \frac{1}{2}\frac{\pi^{-\frac{1-s}{2}}\Gamma(\frac{1-s}{2})}{\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})}\left(1 + i\cot\frac{\pi s}{2}\right)e^{-2\pi iac}a^{s-1} \to \zeta_*(s,0,1)\,.$$

(iii) *As* $(a,c) \to (1^-, 0^+)$,

$$\zeta_*(s,a,c) \quad - \quad \left(c^{-s} + \frac{1}{2}\frac{\pi^{-\frac{1-s}{2}}\Gamma(\frac{1-s}{2})}{\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})}\left(1 - i\cot\frac{\pi s}{2}\right)e^{2\pi i(1-a)c}(1-a)^{s-1}\right) \;\to\; \zeta_*(s,1,0)\,.$$

(iv) *As* $(a,c) \to (0^+, 0^+)$,

$$\zeta_*(s,a,c) \quad - \quad \left(c^{-s} + \frac{1}{2}\frac{\pi^{\frac{1-s}{2}}\Gamma(\frac{1-s}{2})}{\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})}\left(1 + i\cot\frac{\pi s}{2}\right)e^{-2\pi iac}a^{s-1}\right) \;\to\; \zeta_*(s,0,0)\,.$$

Proof. The details are similar to those of Theorem 6.1; we omit them. Note that the limits on the right in cases (i) - (iv) are all $\zeta(s)$, since

$$\zeta_*(s,0,0) = \zeta_*(s,0,1) = \zeta_*(s,1,0) = \zeta_*(s,1,1) = \zeta(s).$$

The limits above are also valid when s is a negative odd integer. ∎

We now deduce Theorem 2.3 using Theorem 6.1 and 6.3.

Proof of Theorem 2.3. We apply Theorems 6.1 and 6.3 for the case $s \in \mathbb{C} \smallsetminus \mathbb{Z}$, and treat remaining case $s \in \mathbb{Z}$ separately.

Suppose that $s \in \mathbb{C} \smallsetminus \mathbb{Z}$. Theorem 6.1(i) gives a continuous extension to the edge $c = 1$, $0 < a' < 1$ for all $s \in \mathbb{C}$. Theorem 6.1(ii) gives a continuous extension to the edge $c = 0$, $0 < a' < 1$ exactly when c^{-s} continuously extends, which is when $\Re(s) < 0$. Theorem 6.1(iii) gives a continuous extension to the edge $a = 1$, $0 < c' < 1$ exactly when $(1-a)^{s-1}$ continuously extends, which is when $\Re(s) > 1$. Theorem 6.1(iv) gives a continuous extension to the edge $a = 0$, $0 < c' < 1$ exactly when a^{s-1} continuously extends, which is when $\Re(s) > 1$. Theorem 6.3 covers the four corners similarly giving a continuous extension to the two corners $(a', c') = (0,1)$ and $(1,1)$ when $\Re(s) > 1$ and in no other cases.

Now suppose that $s = m \in \mathbb{Z}$. If $m \geq 2$ then we are in the region of absolute convergence of the Dirichlet series

$$\zeta(s,a,c) = c^{-s} + \sum_{n=1}^{\infty} e^{2\pi ina}(n+c)^{-s}\,, \tag{6.24}$$

so we get a continuous extension of the sum in the right side of (6.24) to $\square$. The term c^{-s} produces a discontinuity on the side $c = 0$, including the two corners. The case $m \leq -1$ is handled using Lerch's transformation formula (5.12), setting $1 - s = m$. Then $s = |m| + 1$ in (5.12) so the right side has absolutely convergent Dirichlet series, which are continuous except for two terms producing discontinuities along the lines $a = 0$, $a = 1$ and the four corners.

Next suppose that $s = 1$. When $a' = 0$ or 1 the function $\zeta_*(s, a', c)$ has a pole at $s = 1$, so $\zeta_*(s,a,c)$ diverges as $(a,c) \to (0, c')$ or $(1, c')$. For $s = 1$ and $(a,c) \in \square^{\circ}$ the sum in

(6.24) conditionally converges, uniformly in $\epsilon < a < 1-\epsilon$, while the term c^{-s} diverges as $c \to 0^+$. Thus there is no continuous extension to any point on the line $c' = 0$. There is however a continuous extension to the line $c' = 1$, $0 < a' < 1$, by the same reasoning.

The remaining case is $s = 0$. The Lerch transformation formula (5.12) for $s = 1$ expresses $\zeta_*(0, a, c)$ on the lines $a = 0$ and $a = 1$ as a linear combination of $\zeta_*(1, 1-c, a)$ and $\zeta_*(1, c, 1-a)$, with both coefficients nonzero. One of these has a continuous extension to the boundary and the other doesn't, so their sum never does. We conclude from Theorem 6.1(i),(ii) and the holomorphicity of $\zeta_*(s, a, c)$ at $s = 0$ that $\zeta_*(0, a, c)$ has a continuous extension to the line $c = 1$, but not to the line $c = 0$ and not at the four corners of the square. ∎

7. Renormalized Lerch functions

Theorem 6.1 and Theorem 6.3 provide a way to restore continuity on the boundary of the unit square $\square$ for the functions $L^{\pm}(s, a, c)$ by "renormalizing" the functions with correction terms, for $s \in \mathbb{C} \smallsetminus \mathbb{Z}$, as follows. This renormalization is most easily expressed using the following functions.

Definition 7.1. The *Tate gamma functions* $\gamma^{\pm}(s)$, sometimes called *Gelfand-Graev gamma functions*, are given by

$$\gamma^{+}(s) := \frac{\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})}{\pi^{-\frac{1-s}{2}}\Gamma(\frac{1-s}{2})}, \qquad \gamma^{-}(s) := \frac{\pi^{-\frac{s+1}{2}}\Gamma(\frac{s+1}{2})}{\pi^{-\frac{2-s}{2}}\Gamma(\frac{2-s}{2})},$$

These functions were introduced in Tate's thesis [42, p. 317] as normalizing factors in local functional equations at the real place, see also Burnol [7, p. 822]. They satisy the identities

$$\gamma^{\pm}(s)\gamma^{\pm}(1-s) = 1 \text{ for } s \in \mathbb{C}. \tag{7.1}$$

The functions $\gamma^{\pm}(1-s) = \gamma^{\pm}(s)^{-1}$ have a "scattering matrix" interpretation, given in Burnol [6, Sec. 5]. These functions are present in various formulas in §5, where (6.23) can be rewritten

$$\gamma(s) = \frac{1}{2}\left(\gamma^{+}(1-s) - i\gamma^{-}(1-s)\right), \tag{7.2}$$

and the functions $\gamma^{\pm}(1-s)$ appear in formulas of Theorem 6.3.

Definition 7.2. Let $s \in \mathbb{C} \smallsetminus \mathbb{Z}$, and $k = 0$ or 1. Write $(-1)^k = \pm$ and define for $(a, c) \in \square^{\circ}$ the *renormalized Lerch functions*

$$L^{R,\pm}(s, a, c) := L^{\pm}(s, a, c) - S^{\pm}(s, a, c), \tag{7.3}$$

obtained from $L^{\pm}(s, a, c)$ by removing the "correction term"

$$\begin{aligned} S^{\pm}(s, a, c) \;&:=\; \{c^{-s} + (-1)^k e^{-2\pi i a}(1-c)^{-s}\} \\ &+\; i^k \gamma^{\pm}(1-s)\{e^{-2\pi i a c} a^{s-1} + (-1)^k e^{2\pi i(1-a)c}(1-a)^{s-1}\}, \end{aligned} \tag{7.4}$$

in which $\gamma^{\pm}(s)$ denotes the Tate gamma function given in definition 7.1.

The function $S^{\pm}(s,a,c)$ embodies the total contribution of the "bad terms". This definition cannot be further extended to $s \in \mathbb{Z}$, because for each integer s at least one of the gamma factors in $S^{\pm}(s,a,c)$ has a pole. However this definition does further extend in the (a,c)-variables to $(a,c) \in (\mathbb{R} \smallsetminus \mathbb{Z}) \times (\mathbb{R} \smallsetminus \mathbb{Z})$.

Theorem 7.3. (Renormalized Lerch Zeta Functions)

(1) For fixed $s \in \mathbb{C} \smallsetminus \mathbb{Z}$ the "renormalized" functions $L^{R,\pm}(s,a,c)$ continuously extend to the closed unit square $\Box$ in the (a,c) variables.

(2) For fixed $(a,c) \in \Box$ the "renormalized " functions $L^{R,\pm}(s,a,c)$ are meromorphic functions in s having at most simple poles, with all poles in the set $\mathbb{Z}$.

(3) For all $s \in \mathbb{C} \smallsetminus \mathbb{Z}$ and $(a,c) \in \Box$ the completed "renormalized" functions

$$\hat{L}^{R,\pm}(s,a,c) := \pi^{-(\frac{s+k}{2})}\Gamma\left(\frac{s+k}{2}\right)\hat{L}^{R,\pm}(s,a,c)$$

satisfy the functional equations

$$\hat{L}^{R,\pm}(s,a,c) = w_{\pm}e^{-2\pi iac}\hat{L}^{R,\pm}(1-s,1-c,a), \tag{7.5}$$

with $w_+ = 1, w_- = i$.

Remark. This effect of this "renormalization" is to remove the singularities on the boundary of the unit square. However the "renormalized" functions $L^{R,\pm}(s,a,c)$, when extended to $(a,c) \in (\mathbb{R} \smallsetminus \mathbb{Z}) \times (\mathbb{R} \smallsetminus \mathbb{Z})$, still have singularities at all integer values of a or c other than 0 or 1.

Proof. For each fixed $(a,c) \in \Box^{\circ}$ the functions $S^{\pm}(s,a,c)$ are meromorphic functions of s, with simple poles confined to $s \in \mathbb{Z}$. Theorem 2.2 states that $\hat{L}^{\pm}(s,a,c)$ also has this property, with pole set confined to $s \in \{0,1\}$. Thus $L^{R,\pm}(s,a,c)$ inherits this property for $(a,c) \in \Box^{\circ}$.

The functional equation (7.5) for $(a,c) \in \Box^{\circ}$ is inherited from the fact that $\hat{L}^{\pm}(s,a,c)$ satisfies this functional equation (Theorem 2.2), as does the completed "renormalizing" function $\hat{S}^{\pm}(s,a,c) := \pi^{-(\frac{s+k}{2})}\Gamma\left(\frac{s+k}{2}\right)S^{\pm}(s,a,c)$. Indeed one easily verifies the relation

$$\hat{S}^{\pm}(s,a,c) = i^k e^{-2\pi iac}\hat{S}^{\pm}(1-s,1-c,a). \tag{7.6}$$

We now establish that a continuous extension to the boundary of $\Box$ exists for the case $L^{R,+}(s,a,c)$; we omit details for the case $L^{R,-}(s,a,c)$, which is similar. In terms of the expression in Theorems 6.1 and 6.3, we can write

$$L^{R,+}(s,a,c) = A(s,a,c) + B(s,a,c)\,, \tag{7.7}$$

in which

$$\begin{aligned} A(s,a,c) \quad &:= \quad \zeta(s,a,c) - c^{-s} \\ &-\tfrac{1}{2}\gamma^{+}(1-s)\{(1+i\cot\tfrac{\pi s}{2})e^{-2\pi iac}a^{s-1} + (1-i\cot\tfrac{\pi s}{2})e^{2\pi i(1-a)c}(1-a)^{s-1}\}, \end{aligned} \tag{7.8}$$

$$\begin{aligned} B(s,a,c) \quad &:= \quad e^{-2\pi ia}\zeta(s,1-a,1-c) - e^{-2\pi ia}(1-c)^{-s} \\ &-\tfrac{1}{2}\gamma^{+}(1-s)\{(1-i\cot\tfrac{\pi s}{2})e^{-2\pi iac}a^{s-1} + (1+i\cot\tfrac{\pi s}{2})e^{2\pi i(1-a)c}(1-a)^{s-1}\} \end{aligned} \tag{7.9}$$

where $\gamma^+(s) = \frac{\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})}{\pi^{-\frac{1-s}{2}}\Gamma(\frac{s1-}{2})}$. Since $s \notin \mathbb{Z}$ none of the Γ-factors or $\cot\frac{\pi s}{2}$ have poles. In addition $\zeta(s,a,c) = \zeta_*(s,a,c)$ for $(a,c) \in \Box^\circ$, so that Theorem 6.1 applies. We claim that $A(s,a,c)$ continuously extends to the boundary for $s \in \mathbb{C} \smallsetminus \mathbb{Z}$. To see this, when approaching the line $c = 1$, $0 < a' < 1$, the function $\zeta_*(1,a,c)$ continuously extends to it by Theorem 6.1(i), and the other three terms on the right side of (7.8) also continuously extend there. When approaching the line $c = 0$, $0 < a' < 1$, the term $\zeta(s,a,c) - c^{-s}$ continuously extends by Theorem 6.1(ii), while the other two terms on the right side of (7.8) are continuous there. When approaching the line $a = 1$, $0 < c' < 1$, the term $\zeta_*(s,a,c) - \frac{1}{2}\gamma^+(1-s)(1 - i\cot\frac{\pi s}{2})e^{2\pi i(1-a)c}(1-a)^{s-1}$ has a continuous extension by Theorem 6.1(iii) and the other two terms are continuous there. Approaching the line $a = 0$, $0 < c' < 1$, the term $\zeta_*(s,a,c) - \frac{1}{2}\gamma^+(1-s)(1 + i\cot\frac{\pi s}{2})e^{-2\pi iac}a^{s-1}$ has a continuous extension by Theorem 6.1(iv) and the other two terms are continuous there. To get continuity at the four corners of $\Box$, Theorem 6.3 is invoked in a similar fashion. This proves the claim.

In a similar fashion one proves that $B(s,a,c)$ continuously extends to the boundary of $\Box$ when $s \in \mathbb{C} \smallsetminus \mathbb{Z}$. Now (7.7) gives a continuous extension of $L^{R,+}(s,a,c)$ to $\Box$.

The continuous extension to $\Box$ implies that the functional equation (7.5) continues to hold on the boundary $(a,c) \in \partial\Box = \Box \smallsetminus \Box^\circ$. It remains to establish that for $(a,c) \in \partial\Box$ the extended functions $L^{R,\pm}(s,a,c)$ are meromorphic functions of $s \in \mathbb{C}$, having at most simple poles for $s \in \mathbb{Z}$. The crucial fact needed to do this is that the difference functions obtained from subtracting off the "bad terms" are meromorphic in s, which is the content of Lemma 6.2; we omit details. ∎

We now apply Theorem 7.3 to deduce Theorem 2.4, which answers the question: For which values $s \in \mathbb{C}$ do the functions $L^\pm(s,a,c)$ belong to $L^p(\Box, da\, dc)$, for fixed $1 \le p \le 2$?

Proof of Theorem 2.4. We treat first the case of $s \in \mathbb{C} \smallsetminus \mathbb{Z}$. Theorem 7.3 asserts that the functions $L^{R,\pm}(s,a,c)$ are continuous on the unit square (except for integral s), so the property of belonging to $L^p(\Box, dadc)$ for $L^\pm(s,a,c)$ is completely determined by the behavior of the "correction" functions $S^\pm(s,a,c)$, and similarly for linear combinations $F_s(a,c) = c_1L^+(s,a,c) + c_2L^-(s,a,c) \in \mathcal{E}_s$. It then suffices to prove the result for the completed functions $\hat{L}^\pm(s,a,c)$, because the gamma factors take finite nonzero values there.

Suppose first that $1 \le p < 2$. In any linear combination $F_s(a,c)$, one of the pair of correction terms c^{-s} and $e^{-2\pi ia}(1-c)^{-s}$ has a nonzero coefficient, and one of the pair of correction terms $e^{-2\pi iac}a^{s-1}$, and $e^{2\pi i(1-a)c}(1-a)^{s-1}$ has a nonzero coefficient. Furthermore the singularities of each of these terms cannot be cancelled by any linear combination of the other three terms. All functions in both pairs belong to $L^p(\Box, da\, dc)$ for $1 - \frac{1}{p} < \Re(s) < \frac{1}{p}$; thus all linear combinations of $L^\pm(s,a,c)$ belong to $L^p(\Box, da\, dc)$ in this region. Each of the first pair c^{-s} and $e^{-2\pi ia}(1-c)^{-s}$ does not belong to $L^p(\Box, da\, dc)$

for $\Re(s) \ge \frac{1}{p}$, and each of the second pair $e^{-2\pi iac}a^{s-1}$, and $e^{2\pi i(1-a)c}(1-a)^{s-1}$ does not belong to $L^p(\Box, dadc)$ for $\Re(s) \le 1 - \frac{1}{p}$, hence all nonzero linear combinations $F_s(a,c)$ do not belong to $L^p(\Box, da\ dc)$ for $\Re(s) \le 1 - \frac{1}{p}$ and for $\Re(s) \ge \frac{1}{p}$, for non-integral s.

To treat the case $p \ge 2$, it suffices to show the result for $p = 2$, since $L^p(\Box, dadc) \subset L^2(\Box, dadc)$ if $p \ge 2$. We note that each of c^{-s} and $e^{-2\pi ia}(1-c)^{-s}$ does not belong to $L^2(\Box, da\ dc)$ for $\Re(s) \ge \frac{1}{2}$, and each of $e^{-2\pi iac}a^{s-1}$, and $e^{2\pi i(1-a)c}(1-a)^{s-1}$ does not belong to $L^2(\Box, da\ dc)$ for $\Re(s) \le \frac{1}{2}$. As above, we conclude that no nonzero $F_s(a,c)$ belongs to $L^2(\Box, dadc)$, for all $s \in \mathbb{C} \setminus \mathbb{Z}$. Thus assertions (1) and (2) are verified in this case.

It remains to consider the case where $s \in \mathbb{Z}$. We now observe that the Dirichlet series for $L^{\pm}(s,a,c) - |c|^{-s} - (-1)^k e^{-2\pi ia}|1-c|^{-s}$ converges conditionally for $\Re(s) > 0$ and defines a continuous function on $\Box^{\circ} \cup \{(a,0) : 0 < a < 1\} \cup \{(a,1) : 0 < a < 1\}$. As the edges $c = 0$ and $c = 1$ are approached, the growth of the functions $|c|^{-s}$ and $|1-c|^{-s}$ puts the functions $L^{\pm}(s,a,c)$ not in $L^p([\delta, 1-\delta] \times [0,1], da\ dc)$ for any $\delta > 0$ and $\Re(s) \ge 1$. This includes $s = 1, 2, 3, ...$, so $L^{\pm}(s,a,c) \notin L^p(\Box, da\ dc)$ for $s = 1, 2, 3...$ The same argument applies to exclude membership in $L^2(\Box, da\ dc)$ for $s = 1, 2, 3....$ All nonzero linear combinations of these functions are excluded similarly, since it is not possible to cancel both terms $|c|^{-s}$ and $|1-c|^{-s}$ in any linear combination. Finally, a similar argument for $s = 0, -1, -2, ...$ excludes all nonzero linear combinations of $e^{-2\pi iac}L^{\pm}(1-s, 1-c, a)$ from membership in $L^p(\Box, da\ dc)$ for $1 \le p \le 2$, since these functions have a Dirichlet series expansion absolutely convergent in $\Re(s) < 0$ and conditionally convergent in $\Re(s) < 1$, inside $\Box^{\circ}$. ∎

8. Weyl algebra action

The Tate viewpoint on the functional equation uses test functions drawn from the Schwartz space $\mathcal{S} = \mathcal{S}(\mathbb{R})$. The Schwartz space is closed under the action of the *Weyl algebra* $\mathbf{A}_1 := \mathbb{C}[x, \frac{\partial}{\partial x}]$, and carries a (smooth) representation of the Weyl algebra acting as linear operators on $\mathcal{S}$. The Mellin transform intertwines the Weyl algebra action with an action on functions in the s-variable given by difference operators. We use this to define for integer $n \ge 0$ an infinite family $\hat{L}_n(s,a,c)$ of generalized Lerch zeta functions with functional equation, associated to the oscillator representation, whose first two members for $n = 0, 1$ are $\hat{L}^+(s,a,c)$ and $\hat{L}^-(s,a,c)$, respectively.

The *Weyl algebra* is the universal enveloping algebra $U(\mathbf{h}_{\mathbb{R}})$ of the real Heisenberg Lie algebra $\mathbf{h}_{\mathbb{R}}$. Here the *real Heisenberg Lie algebra* is the three-dimensional real Lie algebra

$$\mathbf{h}_{\mathbb{R}} := \mathbb{R}[I, x, \frac{\partial}{\partial x}] = \mathbb{R}[I, D_+, D_-]\ , \tag{8.1}$$

in which

$$D_+ := \sqrt{2\pi}\left(x - \frac{1}{2\pi}\frac{\partial}{\partial x}\right) \quad \text{and} \quad D_- := \sqrt{2\pi}\left(x + \frac{1}{2\pi}\frac{\partial}{\partial x}\right)\ . \tag{8.2}$$

Its commutation relations on the generators I, D_+, D_- are

$$[D_+, D_-] = -2I \quad \text{and} \quad [D_+, I] = [D_-, I] = 0\ . \tag{8.3}$$

The Mellin transforms $\mathcal{M}_k$ acting on even and odd Schwartz functions serve as intertwining operators to carry the Weyl algebra action on functions on $\mathbb{R}^*$ over to function spaces in the s-variable.

Lemma 8.1. *The Weyl algebra* $\mathbf{A}_1$ *acts on the two-sided Mellin transforms* $\mathcal{M}_k$ *of* $f \in \mathcal{S}$*, with* $k = 0, 1 \pmod 2$*, as difference operators in the* s*-variable, as follows:*

$$\mathcal{M}_k(\frac{\partial f}{\partial x})(s) = -(s-1)\mathcal{M}_{k+1}(f)(s-1) \ , \tag{8.4}$$

$$\mathcal{M}_k(xf(x))(s) = \mathcal{M}_{k+1}(f)(s+1) \ . \tag{8.5}$$

As a consequence,

$$\mathcal{M}_k(\frac{\partial^2 f}{\partial x^2})(s) = (s-1)(s-2)\mathcal{M}_k(f)(s-2) \ , \tag{8.6}$$

$$\mathcal{M}_k(x\frac{\partial f}{\partial x})(s) = -s\mathcal{M}_k(f)(s) \ , \tag{8.7}$$

$$\mathcal{M}_k(x^2 f(x))(s) = \mathcal{M}_k(f)(s+2) \ . \tag{8.8}$$

Proof. Integration by parts gives

$$\begin{aligned} \mathcal{M}_k(\frac{\partial f}{\partial x})(s) &= \int_{-\infty}^{\infty} \frac{\partial f}{\partial x}(sgn\, x)^k |x|^s \frac{dx}{|x|} \\ &= -(s-1)\int_{-\infty}^{\infty} f(x)(sgn\, x)^{k+1}|x|^{s-1}\frac{dx}{|x|} \\ &= -(s-1)\mathcal{M}_{k+1}(f)(s-1) \ . \end{aligned} \tag{8.9}$$

The calculation (8.5) is immediate. The remaining three identities follow by substitution. ∎

We consider the Weyl algebra action on the eigenfunctions of the one-dimensional harmonic oscillator Hamiltonian

$$\mathsf{H} := \frac{1}{2}\left(D_+D_- + D_-D_+\right) = -\frac{1}{2\pi}\frac{\partial^2}{\partial x^2} + 2\pi x^2.$$

We work in the infinite-dimensional vector space

$$\mathcal{S}_0 = \mathbb{C}[x]e^{-\pi x^2} \subset \mathcal{S},$$

which is closed under the action of the Weyl algebra. Here

$$\mathcal{S}_0 = \bigoplus_{n \geq 0} \mathbb{C}[\phi_n], \tag{8.10}$$

where the ϕ_n are a basis of eigenfunctions of H, given by $\phi_0(x) = e^{-\pi x^2}$ and

$$\phi_n(x) := D_+^n(\phi_0(x)) = D_+^n(e^{-\pi x^2}), \tag{8.11}$$

The functions $\phi_n(x)$ are explicitly given by

$$\phi_n(x) = 2^{-n/2}H_n(\sqrt{2\pi}x)e^{-\pi x^2}, \tag{8.12}$$

in which $H_n(x)$ are the Hermite polynomials

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n}(e^{-x^2}) \ . \tag{8.13}$$

The functions $\phi_n(x)$ satisfy the (time-independent) Schrödinger equation

$$\mathsf{H}\phi_n = (-\frac{1}{2\pi}\frac{d^2}{dx^2} + 2\pi x^2)\phi_n = (2n+1)\phi_n \ . \tag{8.14}$$

The operators D_+ and D_- are called *raising* and *lowering operators*, due to the relations $D_+\phi_n = \phi_{n+1}, D_-\phi_n = 2n\phi_{n-1}$. The Fourier transform (3.3) acts compatibly with the raising operator

$$\mathcal{F}(D_+f) = -iD_+(\mathcal{F}f), \quad \text{all} \quad f \in \mathcal{S} \ , \tag{8.15}$$

cf. Bump and Ng [5, p. 198]. (Bump and Ng use the convention $\mathcal{F}f(y) = \int_{-\infty}^{\infty} f(x)e^{2\pi ixy}dx$, so that (8.15) differs by a sign change from their equation (1.4).) Since $\phi_0 = \mathcal{F}\phi_0$, we obtain from (8.11) by induction on n that

$$\mathcal{F}\phi_n(x) = (-i)^n\phi_n(x) \ , \quad n \geq 0 \ . \tag{8.16}$$

Bump, Choi, Kurlberg, and Vaaler [4] prove the following results.

Proposition 8.2. *For $n \geq 0$ the Mellin transforms of $\phi_n(x)$ satisfy*

$$\begin{aligned} \mathcal{M}_0(\phi_{2n}(x))(s) &= \pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)p_n(s), && (8.17)\\ \mathcal{M}_1(\phi_{2n+1}(x))(s) &= \sqrt{2\pi}\,\pi^{-\frac{s+1}{2}}\Gamma\left(\frac{s+1}{2}\right)q_n(s), && (8.18) \end{aligned}$$

in which $p_n(s)$ and $q_n(s)$ are real polynomials of degree n. These polynomials satisfy the functional equations

$$\begin{aligned} p_n(1-s) &= (-1)^n p_n(s)\,, \quad n \geq 0,\\ q_n(1-s) &= (-1)^n q_n(s)\,, \quad n \geq 0. \end{aligned} \tag{8.19}$$

The zeros of $p_n(s)$, $q_n(s)$ all lie on the critical line $\Re(s) = 1/2$.

Proof. This is proved in Bump et al. [4, Theorem 1]. The last assertion of this lemma is the "local Riemann hypothesis" studied in Bump and Ng [5], Bump et al [4], Kurlberg [21], and Olofsson[37]. ∎

Lemma 8.1 yields recursion relations for the polynomials $p_n(s)$ and $q_n(s)$, namely

$$\mathcal{M}_k(\phi_{n+1})(s) = \mathcal{M}_k(D_+\phi_n)(s) = \sqrt{2\pi}\mathcal{M}_{k+1}(\phi_n)(s+1) + \frac{1}{\sqrt{2\pi}}(s-1)\mathcal{M}_{k+1}(\phi_n)(s-1), \tag{8.20}$$

where subscripts in $\mathcal{M}_k$ are given (mod 2). From this one readily finds that the polynomials p_n and q_n satisfy $p_0(s) = q_0(s) = 1$ and

$$\begin{aligned} p_{n+1}(s) &= sq_n(s+1) + (s-1)q_n(s-1), && (8.21)\\ q_{n+1}(s) &= p_{n+1}(s+1) + p_{n+1}(s-1)\,. && (8.22) \end{aligned}$$

These in turn imply that $p_n(x), q_n(x) \in \mathbb{Z}[x]$. The first few polynomials $p_n(x)$ and $q_n(x)$ are given in Table 1.

n	$p_n(s)$	$q_n(s)$
0	1	1
1	$2s-1$	$4s-2$
2	$8s^2-8s+6$	$16s^2-16s+22$
3	$32s^3-16s^2+92s-54$	$64s^3-32s^2+376s-140$
4	$128s^4-128s^3+1360s^2-976s+612$	$256s^4-256s^3+4256s^2-2720s+4200$

Table 1. Values of $p_n(s)$ and $q_n(s)$.

From (8.21) and (8.22) we deduce that that these polynomials satisfy the three-term recurrence relations

$$p_{n+1}(s) = sp_n(s+2) + (2s-1)p_n(s) + (s-1)p_n(s-2)\,, \tag{8.23}$$

and

$$q_{n+1}(s) = (s+1)q_n(s+2) + (2s-1)q_n(s) + (s-2)q_n(s-2)\,. \tag{8.24}$$

Three-term recurrence relations are associated to orthogonal polynomials, with respect to a suitable measure, which is explicitly given in the following result.

Proposition 8.3. *(Orthogonal polynomials) For $n \ge 0$ the polynomials*

$$P_n(x) := p_n(\frac{1}{2} + ix) \tag{8.25}$$

are orthogonal polynomials with respect to the measure $|\Gamma(\frac{1}{4} + \frac{ix}{2})|^2 dx$ on the real x-axis. For $n \ge 0$, the polynomials

$$Q_n(x) := q_n(\frac{1}{2} + ix) \tag{8.26}$$

are orthogonal polynomials with respect to the measure $|\Gamma(\frac{3}{4} + \frac{ix}{2})|^2 dx$ on the real x-axis.

Proof. This was observed by Bump et al.[4, pp. 3– 4]. These polynomials are certain Meixner-Pollaczek polynomials $P_n^{(\lambda)}(x;\theta)$, in the notation of Koelink and Swarttouw [20, Sect. 1.7]. Namely,

$$P_n(x) = P_n^{(\frac{1}{4})}(\frac{x}{2};\frac{\pi}{2}) := \frac{(\frac{1}{2})_n}{n!} i^n \ {}_2F_1(-n, \frac{1}{4} + \frac{ix}{2}; \frac{1}{2}; 2),$$

and

$$Q_n(x) = P_n^{(\frac{3}{4})}(\frac{x}{2};\frac{\pi}{2}) := \frac{(\frac{3}{2})_n}{(n!} i^n \ {}_2F_1(-n, \frac{3}{4} + \frac{ix}{2}; \frac{3}{2}; 2).$$

Here the rising factorial $(\lambda)_n := \lambda(\lambda+1)\cdots(\lambda+n-1)$, see also Kutznetsov [22], [23]. ∎

The Mellin transform and the multiplicative averaging operator $A^{a,c}$ given in definition 3.1 map the infinite dimensional vector space $\mathcal{S}_0$ to a vector space $V_{a,c}$ of functions in the s-variable, via

$$\phi(x) \mapsto \mathcal{M}(A^{a,c}[\phi])(s). \tag{8.27}$$

This map is an intertwining map defining a Weyl algebra action on the vector space $V_{a,c}$. The image of the Hermite basis under this intertwining gives a basis of the vector space $V_{a,c}$, which we show below is expressible in terms of Lerch zeta functions.

Definition 8.4. For each $n \in \mathbb{Z}_{\geq 0}$ and fixed $(a,c) \in \Box^{\circ}$ the *generalized (completed) Lerch zeta function* $\hat{L}_n(s,a,c)$ is

$$\begin{aligned} \hat{L}_n(s,a,c) &:= \frac{1}{(2\pi)^{n/2}} \mathcal{M}(2A^{a,c}[\phi_n])(s) \\ &= \frac{2}{(2\pi)^{n/2}} \int_0^{\infty} \left(\sum_{m \in \mathbb{Z}} \phi_n((m+c)x) e^{2\pi i m a} \right) x^{s-1} dx. \end{aligned} \tag{8.28}$$

Note that for $n = 0, 1$ the function $\hat{L}_0(s,a,c) = \hat{L}^+(s,a,c)$ and $\hat{L}_1(s,a,c) = \hat{L}^-(s,a,c)$. The generalized Lerch zeta functions $\hat{L}_n(s,a,c)$ defined using the Hermite basis functions ϕ_n have a simple relation to the functions $L^{\pm}(s,a,c)$, and satisfy similar functional equations, as follows.

Theorem 8.5. *For each* $n \in \mathbb{Z}_{\geq 0}$ *and fixed real* (a,c) *with* $0 < a, c < 1$ *the generalized (completed) Lerch zeta function*

$$\hat{L}_n(s,a,c) := \frac{1}{(2\pi)^{n/2}} \mathcal{M}(2A^{a,c}[\phi_n])(s) = \frac{2}{(2\pi)^{n/2}} \int_0^{\infty} A^{a,c}[\phi_n](x) \; x^{s-1} dx \tag{8.29}$$

extend to entire functions of s*. These functions are given by*

$$\hat{L}_{2n}(s,a,c) = \frac{1}{(2\pi)^{n/2}} p_n(s) \hat{L}^+(s,a,c), \tag{8.30}$$

$$\hat{L}_{2n+1}(s,a,c) = \frac{1}{(2\pi)^{n/2}} \sqrt{2\pi}\, q_n(s) \hat{L}^-(s,a,c) \; . \tag{8.31}$$

For each $n \geq 0$ *they satisfy the functional equation*

$$\hat{L}_n(s,a,c) = i^n \; e^{-2\pi i a c} \hat{L}_n(1-s, 1-c, a) \; . \tag{8.32}$$

Proof. It follows from the definition of $\hat{L}_n(s,a,c)$ and the relation $2A^{a,c} = A_0^{a,c} + A_1^{a,c}$, together with Lemma 3.3, that

$$\hat{L}_n(s,a,c) = \frac{1}{(2\pi)^{n/2}} \{ \mathcal{M}_0(\phi_n)(s) L^+(s,a,c) + \mathcal{M}_1(\phi_n)(s) L^-(s,a,c) \} \; . \tag{8.33}$$

Now $\phi_{2n}(x)$ is an even function and $\phi_{2n+1}(x)$ is an odd function, so that $\mathcal{M}_k(\phi_n)(s) = 0$ if $k \not\equiv n \pmod 2$. Thus we obtain (8.30) and (8.31) according as n is even or odd. The functional equations (8.32) now follow from that of $\hat{L}^{\pm}(s,a,c)$ in Theorem 2.1 and of $p_n(s)$ and $q_n(s)$ in Lemma 8.2. The entire function property of each $\hat{L}_n(s,a,c)$ in the s-variable is inherited from $\hat{L}^{\pm}(s,a,c)$ in Theorem 2.1 and the fact that $p_n(s)$ and $q_n(s)$ are polynomials. ∎

Remarks. (1) Theorem 8.5 shows that the action of the Fourier transform on $\mathcal{S}_0$ carries over an operator of order 4 acting the vector space $V_{a,c}$, which is compatible with the Weyl algebra action.

(2) The results of Bump et al.[4] for the families of polynomials $\{p_n(s)\}$ (resp. $\{q_n(s)\}$) may alternatively be viewed as attached to the Lie algebra $\mathbf{sl}(2)$, viewed as

$$\mathbf{sl}(2) = \mathbb{R}\left[x^2, \frac{1}{2}\left(x\frac{\partial}{\partial x} + \frac{\partial}{\partial x} x \right), \frac{\partial^2}{\partial x^2} \right], \tag{8.34}$$

rather to its universal enveloping algebra $U(\mathbf{sl}(2))$. The universal enveloping algebra $U(\mathbf{sl}(2))$ is a subalgebra of the Weyl algebra $\mathbf{A}_1 = U(\mathbf{h}_{\mathbb{R}})$. As a vector space over $\mathbb{R}$, $U(\mathbf{sl}(2))$ is spanned by all monomials in x and $\frac{\partial}{\partial x}$ of even degree in $\mathbf{A}_1$, counting x and $\frac{\partial}{\partial x}$ as degree one. The members of each family $\{p_n(s)\}$ (resp. $\{q_n(s)\}$) (multiplied by scalar constants) are produced from the bottom element $p_0(s)$ (resp. $q_0(s)$ by repeatedly applying the raising operator D_+^2.

(3) For comparison with the real Heisenberg Lie algebra treated here, the complex Heisenberg algebra $\mathbf{h}_{\mathbb{C}} = \mathbf{h}_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}$ used in quantum mechanics has position operator $p = M_x$ and momentum operator $q = \frac{1}{i}\frac{\partial}{\partial x}$, so that $[p, q] = -iI$. The creation operator $a = p + \frac{i}{2\pi} q$ and annihilation operator $a^* = p - \frac{i}{2\pi} q$ satisfy $[a, a^*] = I$ and $H = \frac{1}{2}(aa^* + a^*a)$.

References

[1] T. M. Apostol, On the Lerch zeta function, *Pacific J. Math.* **1** (1951), 161–167.

[2] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer-Verlag: New York, 1976.

[3] B. Berndt, Two new proofs of Lerch's functional equation, *Proc. Amer. Math. Soc.* **32** (1972), 403–408.

[4] D. Bump, K-K. Choi, P. Kurlberg and J. Vaaler, A Local Riemann Hypothesis, I, Math. Zeitschrift **233** (2000), 1–18.

[5] D. Bump and E. K.-S. Ng, On Riemann's zeta function, Math. Z. **192** (1986) 195–204.

[6] J.-F. Burnol, Des Équations de Dirac et de Schrödinger pour la transformation de Fourier, C. R. Acad. Sci. Paris **336** (2003), 919–924.

[7] J.-F. Burnol, On Fourier and zeta(s), Forum Math. **16** (2004), 789–840.

[8] P. Cartier, Quantum Mechanical Commutation Relations and Theta Functions, in: *Algebraic Groups and Discontinuous Subgroups*, Proc. Symp. Pure Math. Vol. 9, Amer. Math. Soc.: Providence, RI 1966, pp. 361–383.

[9] H. Davenport, *Multiplicative Number Theory*, Second Edition. Revised by H. L. Montgomery. Springer-Verlag: New York 1980.

[10] C. Deninger, *On the Γ-factors attached to motives,* Invent. Math. **104** (1991), 245–261.

[11] C. Deninger, *Local L-factors of motives and regularized determinants*, Invent. Math. **107** 91992), 135–150.

[12] C. Deninger, *Motivic L-functions and regularized determinants*, pp. 707–743 in: *Motives* (U. Jannsen, S. Kleiman, J.-P. Serre, Eds.), Proc. Symp. Pure Math, Vol 55, Part I, American Math. Soc. : Providence RI 1994.

[13] A. Erdélyi et al., *Higher Transcendental Functions – Vol. I*, M. Graw-Hill: New York 1953.

[14] R. Garunkštis, Growth of the Lerch zeta function, Lithuanian Math. J. **45** (2005), 34–43.

[15] R. Garunkštis, J. Steuding, Do Lerch zeta functions satisfy the Lindelöf hypothesis?, in: *Analtyic and probabilistic methods in number theory (Palanga 2001)*, 61–74, TEV: Vilnius 2002.

[16] G. H. Hardy, *Divergent Series*, Oxford University Press: Oxford, 1949.

[17] R. Howe, On the role of the Heisenberg group in harmonic analysis, Bull. Amer. Math. Soc. (New Series) **3** (1980), 821–843.

[18] A. Hurwitz, Einige Eigenshaften der Dirichlet'schen Funktionen $F(s) = \sum \left(\frac{D}{n}\right) \frac{1}{n^s}$, dei bei der Bestimmung der Klessenzahlen binarer quadratischen formen auftreten, *Zeitschrift für Math. und Physik* **27** (1882), 86–101.

[19] S. Kanemitsu and H. Terada, *Vistas of Special Functions*, World Scientific: Singapore 2007.

[20] R. Koekoek and R. F. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue,*, Delft Univ. Techonology, Dept. of Technical Mathematics and Informatics, Report no. 98-17 (1998), 170pp.

[21] P. Kurlberg, A Local Riemann Hypothesis, II, Math. Zeitschrift **233** (2000), 21–37.

[22] A. Kuznetsov, Integral representations for the Dirichlet L-functions and their expansions in Meixner-Pollaczek polynomials and rising factorials, Integral Transforms and Special Functions **18** (2007), No. 11-12, 809–817.

[23] A. Kuznetsov, Expansion of the Riemann Ξ function in Meixner-Pollaczek polynomials, Canadian Math. Bulletin **51** (2008), No. 4, 561–569.

[24] J. C. Lagarias, Number Theory Zeta Functions and Dynamical Zeta Functions, in: T. Branson, Ed. *Spectral Problems in Geometry and Arithmetic, (Iowa City, IA 1997)*, Contemporary Math. Vol. 237, AMS: Providence 1999, pp. 45–86.

[25] J. C. Lagarias, paper in preparation.

[26] J. C. Lagarias and W.-C. W. Li, The Lerch zeta function II. Analytic continuation, preprint.

[27] J. C. Lagarias and W.-C. W. Li, The Lerch zeta function III. Polylogarithms and special values, preprint.

[28] J. C. Lagarias and W.-C. W. Li, The Lerch zeta function IV. Hecke operators, in preparation.

[29] A. Laurenčikas and R. Garunkštis, *The Lerch zeta-function*, Kluwer Academic Publishers: Dordrecht 2002.

[30] M. Lerch, Note sur la fonction $\mathcal{R}(w,x,s)\sum_{k=0}^{\infty}\frac{e^{2k\pi ix}}{(w+k)^s}$, *Acta Math.* **11** (1887), 19–24.

[31] R. Lipschitz, Untersuchung einer aus vier Elementen gebildeten Reihe, *J. Reine Angew.* **54** (1857), 313–328.

[32] R. Lipschitz, Untersuchung der Eigenshaften einer Gattung von undenlichen Reihen, *J. Reine Angew.* **105** (1889), 127–156.

[33] M. Mikolas, New proof and extension of the functional equation of Lerch's zeta function, *Ann. Univ. Sci. Budapest - Eötvos Sect. Math.* **14** (1971), 111–116.

[34] J. Milnor, On polylogarithms, the Hurwitz zeta function, and the Kubert identities, Enseign. Math. **29** (1983), 281–322.

[35] D. Mumford, *Tata Lectures on Theta I*, Birkhäuser: Boston 1983.

[36] E. Oberhettinger, Note on the Lerch zeta function, *Pacific J. Math.* **6** (1956), 117–120.

[37] R. Olofsson, Local Riemann Hypothesis for complex numbers, preprint `math.NT/0605063`.

[38] S. J. Patterson, *An introduction to the theory of the Riemann zeta function*, Cambridge University Press, Cambridge 1988.

[39] D. Ramakrishnan and R. J. Valenza, *Fourier Analysis on Local Fields*, Springer-Verlag: New York 1999.

[40] M. Schröter and C. Soulé, *On a result of Deninger concerning Riemann's zeta function,* pp. 745–747 in: *Motives* (U. Jannsen, S. Kleiman, J.-P. Serre, Eds.), Proc. Symp. Pure Math, Vol 55, Part I, American Math. Soc. : Providence RI 1994.

[41] H. M. Srivastava and J. Choi, *Series Associated with the Zeta and Related Functions*, Kluwer Academic Publishers: Dordrecht 2001.

[42] J. Tate, Fourier Analysis in Number Fields and Hecke's Zeta-Functions, in: *Algebraic Number Theory* (J. W. S. Cassels and A. Fröhlich, Eds.), Academic Press: London 1967, pp. 305–347.

[43] A. Weil, *Elliptic Functions according to Eisenstein and Kronecker*, Springer-Verlag, New York 1976.

Department of Mathematics, University of Michigan, Ann Arbor, MI 48109-1043,USA
E-mail address: `lagarias@umich.edu`

Department of Mathematics, Pennsylvania State University, University Park, PA 16802-8401,USA
E-mail address: `wli@math.psu.edu`